

Europäisches Patentamt
European Patent Office
Office européen des brevets



(11) **EP 0 900 568 A2**

(12)

EUROPEAN PATENT APPLICATION

(43) Date of publication:
10.03.1999 Bulletin 1999/10

(51) Int Cl.⁶ **A61K 31/505**
// (A61K31/505, 31:195, 31:15)

(21) Application number: **98307181.2**

(22) Date of filing: **04.09.1998**

(84) Designated Contracting States:
**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE**
Designated Extension States:
AL LT LV MK RO SI

- **Menniti, Frank Samuel**
Mystic, Connecticut 06355 (US)
- **Welch, Willard KcKowan, Jr.**
Mystic, Connecticut 6355 (US)

(30) Priority: **05.09.1997 US 58098 P**

(71) Applicant: **Pfizer Products Inc.**
Groton, CT 06340-5146 (US)

(74) Representative: **McMunn, Watson Palmer et al**
Pfizer Limited
Patents Department
Ramsgate Road
Sandwich, Kent CT13 9NJ (GB)

(72) Inventors:
• **Chenard, Bertrand Leo**
Waterford, Connecticut 06385 (US)

(54) **AMPA antagonists for the treatment of dyskinesias associated with dopamine agonist therapy**

(57) The invention relates to a method of treating dyskinesias associated with dopamine agonist therapy in a mammal which comprises administering to said mammal a compound, as defined herein, which is an

antagonist of the AMPA receptor. Dopamine agonist therapy, as referred to in the present invention, is generally used in the treatment of a central nervous system disorder such as Parkinson's disease.

EP 0 900 568 A2

Description

Background Of The Invention

[0001] This invention relates to a method of administering AMPA receptor antagonists to treat dyskinesias in mammals, such as humans, resulting from the use of dopamine agonist therapy. Dopamine agonist therapy, as referred to in the present invention, is generally used in the treatment of a central nervous system disorder such as Parkinson's disease. In particular, this invention relates to the treatment of such dyskinesias using one or more AMPA receptor antagonists that are disclosed and claimed in PCT international application number PCT/IB97/00134 (publication no. WO 9743276, PCT/IB98/00150, EP98304319.0, EP98304522.0, PCT/IB98/00151 and United States patent application (provisional no. 60/057990) entitled "Novel Atropisomers Of 2,3-Disubstituted-(5,6)-Heteroaryl-fused-Pyrimidin-4-ones" filed July 23, 1993 with Bertrand L. Chenard and Willard M. Welch named as inventors. The foregoing United States provisional and PCT international patent applications are incorporated herein by reference in their entirety, and copies of the specifications have been placed in the file of the present application.

[0002] Dyskinesias are involuntary physical movements which may include chorea, tremor, ballism, dystonia, athetosis, myoclonus and tic. Dyskinesias often result from treatment of the physical symptoms of Parkinson's disease. Parkinson's disease is characterized by tremor, rigidity, bradykinesia and postural instability. Such motor abnormalities may be reduced by therapies which increase dopamine receptor stimulation. These therapies include drugs which directly stimulate dopamine receptors (such as bromocriptine) or increase the levels of dopamine (such as L-dopa or drugs which inhibit dopamine metabolism). In the present invention, such therapies which increase dopamine receptor stimulation are referred to generally as dopamine agonist therapy. After a period of chronic administration of dopamine agonist therapy to treat Parkinson's disease, new motor abnormalities may emerge. The motor abnormalities associated with dopamine agonist therapy include choreatic dyskinesias and dystonias. The present invention relates to the treatment of dyskinesias associated with dopamine agonist therapy in the treatment of a central nervous system (CNS) disorder, in particular Parkinson's disease, through the administration of an AMPA receptor antagonist as provided below.

[0003] The compounds that may be used in accord with the present invention are antagonists of the AMPA subtype of the glutamate receptor. Glutamate is the principal excitatory neurotransmitter in the central nervous system of mammals. Glutamate synaptic transmission is mediated by several families of receptors including the α -amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid (AMPA), N-methyl-D-aspartate (NMDA), kainic acid (KA), and metabotropic receptors. The AMPA receptor subtype mediates fast excitatory transmission throughout the brain, including areas involved in movement. By inhibiting the AMPA receptor through administration of an AMPA receptor antagonist, dyskinesias associated with dopamine agonist therapy may be treated in accord with the present invention as described below.

[0004] AMPA receptor antagonists are referred to in several published patents including the following issued United States patents (listed by patent number followed by issue date in parentheses): 5,654,303 (August 5, 1997); 5,639,751 (June 17, 1997); 5,614,532 (March 25, 1997); 5,614,508 (March 25, 1997); 5,606,062 (February 25, 1997); 5,580,877 (December 3, 1996); 5,559,125 (September 24, 1996); 5,559,106 (September 24, 1996); 5,532,236 (July 2, 1996); 5,527,810 (June 18, 1996); 5,521,174 (May 28, 1996); 5,519,019 (May 21, 1996); 5,514,680 (May 7, 1996); 5,631,373 (May 20, 1997); 5,622,952 (April 22, 1997); 5,620,979 (April 15, 1997); 5,510,338 (April 23, 1996); 5,504,085 (April 2, 1996); 5,475,008 (December 12, 1995); 5,446,051 (August 29, 1995); 5,426,106 (June 20, 1995); 5,420,155 (May 30, 1995); 5,407,935 (April 18, 1995); 5,399,696 (March 21, 1995); 5,395,827 (March 7, 1995); 5,376,748 (December 27, 1994); 5,364,876 (November 15, 1994); 5,356,902 (October 18, 1994); 5,342,946 (August 30, 1994); 5,268,378 (December 7, 1993); and 5,252,584 (October 12, 1993).

Summary Of The Invention

[0005] This invention relates to the use of a compound selected from groups (A), (B), (C), (D), (E), or (F) or a pharmaceutically acceptable salt thereof, in the manufacture of a medicament for treating dyskinesias associated with dopamine agonist therapy in a mammal, such as a human, wherein groups (A), (B), (C), (D), (E), and (F) are defined as follows:

(A) (S)-3-(2-chloro-phenyl)-2-[2-(5-diethylaminomethyl-2-fluoro-phenyl)-vinyl]-6-fluoro-3H-quinazolin-4-one;

(S)-3-(2-chloro-phenyl)-2-[2-(6-diethylaminomethyl-pyridin-2-yl)-vinyl]-6-fluoro-3H-quinazolin-4-one;

(S)-3-(2-chloro-phenyl)-2-[2-(4-diethylaminomethyl-pyridin-2-yl)-vinyl]-6-fluoro-3H-quinazolin-4-one;

(S)-3-(2-chloro-phenyl)-2-[2-(6-ethylaminomethyl-pyridin-2-yl)-vinyl]-6-fluoro-3H-quinazolin-4-one;

(S)-3-(2-bromo-phenyl)-2-[2-(6-diethylaminomethyl-pyridin-2-yl)-vinyl]-6-fluoro-3H-quinazolin-4-one;

(S)-3-(2-chloro-phenyl)-6-fluoro-2-[2-(6-methoxymethyl-pyridin-2-yl)-vinyl]-3H-quinazolin-4-one;

(S)-3-(2-chloro-phenyl)-6-fluoro-2-[2-(4-methyl-pyrimidine-2-yl)-vinyl]-3H-quinazolin-4-one;
 (S)-3-(2-chloro-phenyl)-6-fluoro-2-[2-[6-(isopropylamino-methyl)-pyridin-2-yl]-ethyl]-3H-quinazolin-4-one;
 (S)-6-fluoro-2-[2-(2-methyl-thiazol-4-yl)-vinyl]-3-(2-methyl-phenyl)-3H-quinazolin-4-one;
 (S)-3-(2-chloro-phenyl)-6-fluoro-2-[2-(2-methyl-thiazol-4-yl)-vinyl]-3H-quinazolin-4-one;
 5 (S)-2-[2-(2-dimethylaminomethyl-thiazol-4-yl)-vinyl]-6-fluoro-3-(2-fluoro-phenyl)-3H-quinazolin-4-one;
 (S)-3-(2-bromo-phenyl)-6-fluoro-2-[2-(2-methyl-thiazol-4-yl)-vinyl]-3H-quinazolin-4-one;
 (S)-3-(2-chloro-phenyl)-2-[2-(2-methyl-thiazol-4-yl)-vinyl]-3H-quinazolin-4-one;
 (S)-3-(2-chloro-phenyl)-6-fluoro-2-(2-pyridin-2-yl-vinyl)-3H-quinazolin-4-one;
 10 (S)-3-(2-bromo-phenyl)-2-(2-pyridin-2-yl-vinyl)-3H-quinazolin-4-one;
 (S)-6-chloro-2-(2-pyridin-2-yl-vinyl)-3-o-tolyl-3H-quinazolin-4-one;
 (S)-3-(2-chloro-phenyl)-2-[2-(6-methyl-pyridin-2-yl)-vinyl]-3H-quinazolin-4-one;
 (S)-6-chloro-2-[2-(6-methyl-pyridin-2-yl)-vinyl]-3-o-tolyl-3H-quinazolin-4-one;
 (S)-3-(2-chloro-phenyl)-6-fluoro-2-(2-pyridin-2-yl-ethyl)-3H-quinazolin-4-one;
 15 (S)-6-[2-[3-(2-chloro-phenyl)-6-fluoro-4-oxo-3,4-dihydro-quinazolin-2-yl]-vinyl]-pyridine-2-carbaldehyde;
 (S)-3-(2-chloro-phenyl)-6-fluoro-2-[2-(6-methylaminomethyl-pyridin-2-yl)-vinyl]-3H-quinazolin-4-one;
 (S)-N-(6-[2-[3-(2-chloro-phenyl)-6-fluoro-4-oxo-3,4-dihydro-quinazolin-2-yl]-vinyl]-pyridin-2-ylmethyl)-N-methyl-acetamide;
 (S)-6-[2-[3-(2-chloro-phenyl)-6-fluoro-4-oxo-3,4-dihydro-quinazolin-2-yl]-vinyl]-pyridine-2-carbonitrile;
 20 (S)-3-(2-fluoro-phenyl)-2-(2-pyridin-2-yl-vinyl)-3H-quinazolin-4-one;
 (S)-3-(2-bromo-phenyl)-6-fluoro-2-(2-pyridin-2-yl-vinyl)-3H-quinazolin-4-one;
 (S)-3-(4-bromo-2-chloro-phenyl)-6-fluoro-2-(2-pyridin-2-yl-vinyl)-3H-quinazolin-4-one;
 (S)-3-(2-chloro-phenyl)-2-[2-(6-diethylaminomethyl-pyridin-2-yl)-vinyl]-3H-quinazolin-4-one;
 (S)-N-(6-[2-[3-(2-chloro-phenyl)-6-fluoro-4-oxo-3,4-dihydro-quinazolin-2-yl]-vinyl]-pyridin-2-ylmethyl)-N-ethyl-acetamide;
 25 (S)-3-(2-chloro-phenyl)-6-fluoro-2-[2-(6-fluoromethyl-pyridin-2-yl)-vinyl]-3H-quinazolin-4-one;
 (S)-3-(2-chloro-phenyl)-6-fluoro-2-[2-(6-pyrrolidin-1-ylmethyl-pyridin-2-yl)-ethyl]-3H-quinazolin-4-one;
 (S)-3-(2-chloro-phenyl)-2-[2-(6-[[ethyl-(2-hydroxy-ethyl)-amino]-methyl]-pyridin-2-yl)-vinyl]-6-fluoro-3H-quinazolin-4-one;
 30 (S)-3-(2-chloro-phenyl)-6-fluoro-2-[2-[6-(isopropylamino-methyl)-pyridin-2-yl]-vinyl]-3H-quinazolin-4-one;
 (S)-3-(2-chloro-phenyl)-6-fluoro-2-[2-[6-(2-methyl-piperidin-1-ylmethyl)-pyridin-2-yl]-vinyl]-3H-quinazolin-4-one;
 (S)-3-(2-chloro-phenyl)-2-[2-(6-ethoxymethyl-pyridin-2-yl)-vinyl]-6-fluoro-3H-quinazolin-4-one;
 (S)-3-(2-chloro-phenyl)-2-[2-[6-(2,5-dihydro-pyrrol-1-ylmethyl)-pyridin-2-yl]-vinyl]-6-fluoro-3H-quinazolin-4-one;
 35 (S)-3-(2-chloro-phenyl)-6-fluoro-2-[2-[6-(4-methyl-piperidin-1-ylmethyl)-pyridin-2-yl]-vinyl]-3H-quinazolin-4-one;
 (S)-6-bromo-2-[2-(6-methyl-pyridin-2-yl)-vinyl]-3-o-tolyl-3H-quinazolin-4-one;
 (S)-6-bromo-2-(2-pyridin-2-yl-vinyl)-3-o-tolyl-3H-quinazolin-4-one;
 40 (S)-6-fluoro-3-(2-fluoro-phenyl)-2-(2-pyridin-2-yl-vinyl)-3H-quinazolin-4-one;
 (S)-3-(2-chloro-phenyl)-6-methyl-2-(2-pyridin-2-yl-vinyl)-3H-quinazolin-4-one;
 (S)-3-(2-chloro-phenyl)-2-[2-(6-dimethylaminomethyl-pyridin-2-yl)-vinyl]-6-fluoro-3H-quinazolin-4-one;
 (S)-6-fluoro-3-(2-fluoro-phenyl)-2-[2-(6-methyl-pyridin-2-yl)-vinyl]-3H-quinazolin-4-one;
 (S)-3-(2-chloro-phenyl)-2-[2-(6-[[2-dimethylamino-ethyl]-methyl-amino]-methyl)-pyridin-2-yl)-vinyl]-6-fluoro-3H-quinazolin-4-one;
 45 (S)-3-(2-chloro-phenyl)-6-fluoro-2-[2-(6-hydroxymethyl-pyridin-2-yl)-vinyl]-3H-quinazolin-4-one;
 (S)-acetic acid 6-[2-[3-(2-chloro-phenyl)-6-fluoro-4-oxo-3,4-dihydro-quinazolin-2-yl]-vinyl]-pyridin-2-yl methyl ester;
 (S)-6-[2-[3-(2-bromo-phenyl)-6-fluoro-4-oxo-3,4-dihydro-quinazolin-2-yl]-vinyl]-pyridine-2-carbaldehyde;
 50 (S)-3-(2-bromo-phenyl)-2-[2-(6-diethylaminomethyl-pyridin-2-yl)-vinyl]-3H-quinazolin-4-one;
 (S)-acetic acid 6-[2-[3-(2-bromo-phenyl)-6-fluoro-4-oxo-3,4-dihydro-quinazolin-2-yl]-vinyl]-pyridin-2-ylmethyl ester;
 (S)-diethylamino-acetic acid 6-[2-[3-(2-chloro-phenyl)-6-fluoro-4-oxo-3,4-dihydro-quinazolin-2-yl]-vinyl]-pyridin-2-ylmethyl ester;
 55 (S)-3-(2-chloro-phenyl)-2-[2-(6-difluoromethyl-pyridin-2-yl)-vinyl]-6-fluoro-3H-quinazolin-4-one;
 (S)-3-(2-chloro-phenyl)-6-fluoro-2-[2-(6-methoxy-pyridin-2-yl)-vinyl]-3H-quinazolin-4-one;
 (S)-2-[2-[3-(2-chloro-phenyl)-6-fluoro-4-oxo-3,4-dihydro-quinazolin-2-yl]-vinyl]-6-methyl-nicotinonitrile;
 (S)-2-[2-[3-(2-chloro-phenyl)-6-fluoro-4-oxo-3,4-dihydro-quinazolin-2-yl]-ethyl]-6-methyl-nicotinonitrile;
 (S)-3-(2-chloro-phenyl)-6-fluoro-2-(2-pyrimidine-2-yl-ethyl)-3H-quinazolin-4-one;

(S)-3-(2-chloro-phenyl)-2-[2-(4,6-dimethyl-pyrimidine-2-yl)-vinyl]-6-fluoro-3H-quinazolin-4-one;
 (S)-2-[2-[3-(2-chloro-phenyl)-6-fluoro-4-oxo-3,4-dihydro-quinazolin-2-yl]-vinyl]-nicotinonitrile;
 (S)-3-[2-chloro-phenyl]-6-fluoro-2-[2-[6-[(3-methyl-butylamino)-methyl]-pyridin-2-yl]-ethyl]-3H-quinazolin-4-one;
 (S)-2-[2-[3-(2-chloro-phenyl)-6-fluoro-4-oxo-3,4-dihydro-quinazolin-2-yl]-ethyl]-nicotinonitrile;
 (S)-2-[2-[6-chloro-4-oxo-3-o-tolyl-3,4-dihydro-quinazolin-2-yl]-vinyl]-benzonitrile;
 (S)-2-[2-[3-(2-chloro-phenyl)-6-fluoro-4-oxo-3,4-dihydro-quinazolin-2-yl]-vinyl]-4-methyl-benzonitrile;
 (S)-3-(2-bromo-phenyl)-6-fluoro-2-[2-(6-hydroxymethyl-pyridin-2-yl)-vinyl]-3H-quinazolin-4-one; and
 (S)-3-(2-chloro-phenyl)-6-fluoro-2-[2-(6-pyrrolidin-1-ylmethyl-pyridin-2-yl)-vinyl]-3H-quinazolin-4-one;

(B) (S)-6-fluoro-2-[2-(2-fluoro-phenyl)-vinyl]-3-(2-methyl-pyridin-3-yl)-3H-quinazolin-4-one;

(S)-2-[2-[6-fluoro-3-(2-methyl-pyridin-3-yl)-4-oxo-3,4-dihydro-quinazolin-2-yl]-vinyl]-benzonitrile;
 (S)-2-[2-[6-fluoro-3-(2-methylpyridin-3-yl)-4-oxo-3,4-dihydroquinazolin-2-yl]-vinyl]-benzonitrile;
 (S)-2-[2-[3-(2-chloro-pyridin-3-yl)-6-fluoro-4-oxo-3,4-dihydroquinazolin-2-yl]-vinyl]-benzonitrile;
 (S)-2-[2-[6-fluoro-3-(2-methyl-pyridin-3-yl)-4-oxo-3,4-dihydro-quinazolin-2-yl]-vinyl]-4-methyl-benzonitrile;
 (S)-2-[2-[3-(2-methyl-pyridin-3-yl)-4-oxo-3,4-dihydro-quinazolin-2-yl]-vinyl]-benzonitrile;
 (S)-6-fluoro-3-(2-methyl-pyridin-3-yl)-2-[2-(thiazol-2-yl)-vinyl]-3H-quinazolin-4-one;
 (S)-6-fluoro-3-(2-methyl-pyridin-3-yl)-2-[2-(2-methyl-thiazol-4-yl)-vinyl]-3H-quinazolin-4-one;
 (S)-6-fluoro-3-(2-methyl-pyridin-3-yl)-2-[2-(4-methyl-thiazol-2-yl)-vinyl]-3H-quinazolin-4-one;
 (S)-2-[2-(5-diethylaminomethyl-2-fluoro-phenyl)-vinyl]-6-fluoro-3-(2-methyl-pyridin-3-yl)-3H-quinazolin-4-one;
 (S)-6-fluoro-2-[2-(2-fluoro-5-pyrrolidin-1-ylmethyl-phenyl)-vinyl]-3-(2-methyl-pyridin-3-yl)-3H-quinazolin-4-one;
 (S)-3-(2-chloro-pyridin-3-yl)-2-[2-(2-fluoro-phenyl)-vinyl]-3H-quinazolin-4-one;
 (S)-3-(2-chloro-pyridin-3-yl)-6-fluoro-2-[2-(6-methyl-phenyl-2-yl)-vinyl]-3H-quinazolin-4-one;
 (S)-3-(2-chloro-pyridin-3-yl)-6-fluoro-2-[2-(fluoro-phenyl)-vinyl]-3H-quinazolin-4-one;
 (S)-6-chloro-2-[2-(2-fluoro-phenyl)-vinyl]-3-(2-methyl-pyridin-3-yl)-3H-quinazolin-4-one;
 (S)-6-chloro-2-[2-(2-fluoro-phenyl)-vinyl]-3-(3-methyl-1-oxy-pyridin-4-yl)-3H-quinazolin-4-one;
 (S)-3-[2-(3-(2-chloro-pyridin-3-yl)-6-fluoro-4-oxo-3,4-dihydro-quinazolin-2-yl)-vinyl]-benzaldehyde;
 (S)-3-[2-[3-(2-chloro-pyridin-3-yl)-4-oxo-3,4-dihydro-quinazolin-2-yl]-vinyl]-benzaldehyde;
 (S)-3-(2-chloro-pyridin-3-yl)-6-fluoro-2-[2-(3-hydroxymethyl-phenyl)-vinyl]-3H-quinazolin-4-one;
 (S)-3-(2-chloro-pyridin-3-yl)-2-[2-[3-(1,4-dioxo-8-aza-spiro[4.5]dec-8-ylmethyl)-phenyl]-vinyl]-6-fluoro-3H-quinazolin-4-one;
 (S)-3-(2-chloro-pyridin-3-yl)-6-fluoro-2-[2-[3-(4-pyrrolidin-1-yl-piperidin-1-ylmethyl)-phenyl]-vinyl]-3H-quinazolin-4-one;
 (S)-2-[2-[3-(2-chloro-pyridin-3-yl)-6-fluoro-4-oxo-3,4-dihydro-quinazolin-2-yl]-vinyl]-benzonitrile;
 (S)-2-[2-[3-(2-chloro-pyridin-3-yl)-4-oxo-3,4-dihydro-quinazolin-2-yl]-vinyl]-benzonitrile;
 (S)-2-[2-(2-fluoro-phenyl)-vinyl]-3-(2-methyl-pyridin-3-yl)-3H-quinazolin-4-one;
 (S)-6-fluoro-3-(2-methyl-pyridin-3-yl)-6-fluoro-2-[2-(hydroxy-phenyl)-vinyl]-3H-quinazolin-4-one;
 (S)-6-fluoro-3-(2-methyl-pyridin-3-yl)-2-[2-(2-methyl-thiazol-4-yl)-ethyl]-3H-quinazolin-4-one;
 (S)-6-fluoro-3-(2-chloro-pyridin-3-yl)-2-[2-(2-dimethylamino-methylthiazol-4-yl)-vinyl]-3H-quinazolin-4-one;
 (S)-2-[2-(5-Diethylaminomethyl-2-fluoro-phenyl)-vinyl]-6-fluoro-3-(4-methyl-pyridin-3-yl)-3H-quinazolin-4-one;
 (S)-4-Diethylaminomethyl-2-[2-[6-fluoro-3-(4-methyl-pyridin-3-yl)-4-oxo-3,4-dihydro-quinazolin-2-yl]-vinyl]-benzonitrile;
 (S)-2-[2-(5-Diethylaminomethyl-2-fluoro-phenyl)-vinyl]-6-fluoro-3-(3-methyl-pyrazin-2-yl)-3H-quinazolin-4-one;
 (S)-6-fluoro-3-(2-methyl-pyridin-3-yl)-2-[2-(2-dimethylamino-methylthiazol-4-yl)-vinyl]-3H-quinazolin-4-one;
 (S)-6-fluoro-3-(2-methyl-pyridin-3-yl)-2-[2-(2-methyl-oxazol-4-yl)-vinyl]-3H-quinazolin-4-one;
 (S)-6-fluoro-3-(2-chloro-pyridin-3-yl)-2-[2-(thiazol-2-yl)-vinyl]-3H-quinazolin-4-one;
 (S)-6-fluoro-3-(4-methyl-pyridin-3-yl)-2-[2-(4-methyl-thiazol-2-yl)-vinyl]-3H-quinazolin-4-one;
 (S)-3-(2-chloro-pyridin-3-yl)-6-fluoro-2-[2-(2-hydroxy-phenyl)-vinyl]-3H-quinazolin-4-one; and
 (S)-6-fluoro-2-[2-(2-fluoro-5-pyrrolidin-1-ylmethyl-phenyl)-ethyl]-3-(2-methyl-pyridin-3-yl)-3H-quinazolin-4-one;

(C) 3-(2-chloro-phenyl)-6-fluoro-2-(2-pyridin-2-yl-vinyl)-3H-quinazolin-4-one;

3-(2-bromo-phenyl)-2-(2-pyridin-2-yl-vinyl)-3H-quinazolin-4-one;
 6-chloro-2-(2-pyridin-2-yl-vinyl)-3-o-tolyl-3H-quinazolin-4-one;
 3-(2-chloro-phenyl)-2-[2-(6-methyl-pyridin-2-yl)-vinyl]-3H-quinazolin-4-one;
 6-chloro-2-[2-(6-methyl-pyridin-2-yl)-vinyl]-3-o-tolyl-3H-quinazolin-4-one;
 3-(2-chloro-phenyl)-6-fluoro-2-(2-pyridin-2-yl-ethyl)-3H-quinazolin-4-one;
 6-[2-[3-(2-chloro-phenyl)-6-fluoro-4-oxo-3,4-dihydro-quinazolin-2-yl]-vinyl]-pyridine-2-carbaldehyde;
 3-(2-chloro-phenyl)-6-fluoro-2-[2-(6-methylaminomethyl-pyridin-2-yl)-vinyl]-3H-quinazolin-4-one;
 N-(6-[2-[3-(2-chloro-phenyl)-6-fluoro-4-oxo-3,4-dihydro-quinazolin-2-yl]-vinyl]-pyridin-2-ylmethyl)-N-methyl-
 acetamide;
 3-(2-chloro-phenyl)-2-[2-(4-diethylaminomethyl-pyridin-2-yl)-vinyl]-6-fluoro-3H-quinazolin-4-one;
 6-[2-[3-(2-chloro-phenyl)-6-fluoro-4-oxo-3,4-dihydro-quinazolin-2-yl]-vinyl]-pyridine-2-carbonitrile;
 3-(2-fluoro-phenyl)-2-(2-pyridin-2-yl-vinyl)-3H-quinazolin-4-one;
 3-(2-bromo-phenyl)-6-fluoro-2-(2-pyridin-2-yl-vinyl)-3H-quinazolin-4-one;
 3-(4-bromo-2-chloro-phenyl)-6-fluoro-2-(2-pyridin-2-yl-vinyl)-3H-quinazolin-4-one;
 3-(2-chloro-phenyl)-2-[2-(6-diethylaminomethyl-pyridin-2-yl)-vinyl]-3H-quinazolin-4-one;
 N-(6-[2-[3-(2-chloro-phenyl)-6-fluoro-4-oxo-3,4-dihydro-quinazolin-2-yl]-vinyl]-pyridin-2-ylmethyl)-N-ethyl-
 acetamide;
 3-(2-chloro-phenyl)-6-fluoro-2-[2-(6-fluoromethyl-pyridin-2-yl)-vinyl]-3H-quinazolin-4-one;
 3-(2-chloro-phenyl)-6-fluoro-2-[2-(6-pyrrolidin-1-ylmethyl-pyridin-2-yl)-ethyl]-3H-quinazolin-4-one;
 3-(2-chloro-phenyl)-2-[2-(6-[[ethyl-(2-hydroxy-ethyl)-amino]-methyl]-pyridin-2-yl)-vinyl]-6-fluoro-3H-quinazo-
 lin-4-one;
 3-(2-chloro-phenyl)-6-fluoro-2-[2-[6-(isopropylamino-methyl)-pyridin-2-yl]-vinyl]-3H-quinazolin-4-one;
 3-(2-chloro-phenyl)-6-fluoro-2-[2-[6-(2-methyl-piperidin-1-ylmethyl)-pyridin-2-yl]-vinyl]-3H-quinazolin-4-one;
 3-(2-chloro-phenyl)-2-[2-(6-ethylaminomethyl-pyridin-2-yl)-vinyl]-6-fluoro-3H-quinazolin-4-one;
 3-(2-chloro-phenyl)-2-[2-(6-ethoxymethyl-pyridin-2-yl)-vinyl]-6-fluoro-3H-quinazolin-4-one;
 3-(2-chloro-phenyl)-2-[2-[6-(2,5-dihydro-pyrrol-1-ylmethyl)-pyridin-2-yl]-vinyl]-6-fluoro-3H-quinazolin-4-one;
 3-(2-chloro-phenyl)-6-fluoro-2-[2-[6-(4-methyl-piperidin-1-ylmethyl)-pyridin-2-yl]-vinyl]-3H-quinazolin-4-one;
 6-bromo-2-[2-(6-methyl-pyridin-2-yl)-vinyl]-3-o-tolyl-3H-quinazolin-4-one;
 6-bromo-2-(2-pyridin-2-yl-vinyl)-3-o-tolyl-3H-quinazolin-4-one;
 6-fluoro-3-(2-fluoro-phenyl)-2-(2-pyridin-2-yl-vinyl)-3H-quinazolin-4-one;
 1-benzyl-5-(2-methyl-[1,3]dioxolan-2-yl)-2-oxo-2, 3-dihydro-1H-indole-3-carboxylic acid (3-phenylcarbamoyl-
 phenyl)-amide;
 3-(2-chloro-phenyl)-6-methyl-2-(2-pyridin-2-yl-vinyl)-3H-quinazolin-4-one;
 3-(2-chloro-phenyl)-2-[2-(6-dimethylaminomethyl-pyridin-2-yl)-vinyl]-6-fluoro-3H-quinazolin-4-one;
 6-fluoro-3-(2-fluoro-phenyl)-2-[2-(6-methyl-pyridin-2-yl)-vinyl]-3H-quinazolin-4-one;
 3-(2-chloro-phenyl)-2-[2-(6-[[2-(dimethylamino-ethyl)-methyl-amino]-methyl]-pyridin-2-yl)-vinyl]-6-fluoro-3H-
 quinazolin-4-one;
 3-(2-chloro-phenyl)-6-fluoro-2-[2-(6-hydroxymethyl-pyridin-2-yl)-vinyl]-3H-quinazolin-4-one;
 acetic acid 6-[2-[3-(2-chloro-phenyl)-6-fluoro-4-oxo-3,4-dihydro-quinazolin-2-yl]-vinyl]-pyridin-2-ylmethyl es-
 ter;
 6-[2-[3-(2-bromo-phenyl)-6-fluoro-4-oxo-3,4-dihydro-quinazolin-2-yl]-vinyl]-pyridine-2-carbaldehyde;
 3-(2-bromo-phenyl)-2-[2-(6-diethylaminomethyl-pyridin-2-yl)-vinyl]-6-fluoro-3H-quinazolin-4-one;
 3-(2-bromo-phenyl)-2-[2-(6-diethylaminomethyl-pyridin-2-yl)-vinyl]-3H-quinazolin-4-one;
 acetic acid 6-[2-[3-(2-bromo-phenyl)-6-fluoro-4-oxo-3,4-dihydro-quinazolin-2-yl]-vinyl]-pyridin-2-ylmethyl es-
 ter;
 3-(2-chloro-phenyl)-6-fluoro-2-[2-(6-methoxymethyl-pyridin-2-yl)-vinyl]-3H-quinazolin-4-one;
 diethylamino-acetic acid 6-[2-[3-(2-chloro-phenyl)-6-fluoro-4-oxo-3,4-dihydro-quinazolin-2-yl]-vinyl]-pyridin-
 2-ylmethyl ester;
 6-fluoro-3-(2-methyl-pyridin-3-yl)-2-[2-(2-methyl-thiazol-4-yl)-vinyl]-3H-quinazolin-4-one;
 3-(2-bromo-phenyl)-6-fluoro-2-[2-(6-hydroxymethyl-pyridin-2-yl)-vinyl]-3H-quinazolin-4-one; and,
 3-(2-chloro-phenyl)-6-fluoro-2-[2-(6-pyrrolidin-1-ylmethyl-pyridin-2-yl)-vinyl]-3H-quinazolin-4-one;

(D) 3-(2-chloro-phenyl)-2-[2-(6-diethylaminomethyl)-pyridin-2-yl]-2-hydroxy-vinyl]-6-fluoro-3H-quinazolin-4-one


6-Chloro-3-(2-chloro-phenyl)-2-[2-hydroxy-2-(6-methyl-pyridin-2-yl)-vinyl]-3H-quinazolin-4-one;
 2-[2-[3-(2-Chloro-phenyl)-4-oxo-3,4-dihydro-quinazolin-2-yl]-1-hydroxy-vinyl]-nicotinonitrile;
 2-[2-[3-(2-Chloro-pyrid-3-yl)-6-fluoro-4-oxo-3,4-dihydro-quinazolin-2-yl]-1-hydroxy-vinyl]-nicotinonitrile;
 2-[2-[6-Chloro-3-(2-methyl-phenyl)-4-oxo-3,4-dihydro-quinazolin-2-yl]-1-hydroxy-vinyl]-nicotinonitrile;

3-(2-Chloro-phenyl)-2-[2-(3-diethylaminomethyl-phenyl)-2-hydroxy-ethyl]-6-fluoro-3H-quinazolin-4-one;
 3-(2-Chloro-phenyl)-6-fluoro-2-[2-(3-pyrrolidin-1-ylmethyl-phenyl)-2-hydroxy-ethyl]-3H-quinazolin-4-one;
 3-(2-Chloro-pyrid-3-yl)-2-[2-(3-diethylaminomethyl-phenyl)-2-hydroxy-ethyl]-6-fluoro-3-(2-fluoro-phenyl)-3H-quinazolin-4-one;
 2-[2-(3-Diethylaminomethyl-phenyl)-2-hydroxy-ethyl]-3-(2-fluoro-phenyl)-3H-quinazolin-4-one;
 3-(2-chloro-phenyl)-2-[2-(6-diethylaminomethyl)-pyridin-2-yl]-2-hydroxy-vinyl]-6-fluoro-3H-quinazolin-4-one;
 2-[2-[3-(2-Chloro-pyrid-3-yl)-6-fluoro-4-oxo-3,4-dihydro-quinazolin-2-yl]-1-hydroxy-vinyl]-6-methyl-nicotinon-
 itrile;
 2-[2-[3-(2-Chloro-phenyl)-4-oxo-3,4-dihydro-quinazolin-2-yl]-1-hydroxy-vinyl]-6-methyl-nicotinonitrile;
 2-[2-[6-Chloro-3-(2-chloro-phenyl)-4-oxo-3,4-dihydro-quinazolin-2-yl]-1-hydroxy-vinyl]-6-methyl-nicotinoni-
 trile;
 2-[2-[3-(2-Chloro-phenyl)-6-fluoro-4-oxo-3,4-dihydro-quinazolin-2-yl]-1-hydroxy-vinyl]-6-fluoro-nicotinoni-
 trile;
 2-[2-[3-(2-Chloro-phenyl)-6-fluoro-4-oxo-3,4-dihydro-quinazolin-2-yl]-1-hydroxy-vinyl]-4-fluoro-benzonitrile;
 2-[2-[3-(2-Chloro-phenyl)-6-fluoro-4-oxo-3,4-dihydro-quinazolin-2-yl]-1-hydroxy-vinyl]-4-methyl-benzonitrile;
 2-[2-[3-(2-Chloro-phenyl)-4-oxo-3,4-dihydro-thieno[3,2-d]pyrimidin-2-yl]-1-hydroxy-vinyl]-6-methyl-nicotinon-
 itrile;
 2-[2-[3-(2-methyl-phenyl)-4-oxo-3,4-dihydro-thieno[3,2-d]pyrimidin-2-yl]-1-hydroxy-vinyl]-6-methyl-nicotinon-
 itrile;
 2-[2-[3-(2-Chloro-pyrid-3-yl)-4-oxo-3,4-dihydro-thieno[3,2-d]pyrimidin-2-yl]-1-hydroxy-vinyl]-4-methyl-ben-
 zonitrile;
 2-[2-[3-(2-Chloro-phenyl)-4-oxo-3,4-dihydro-thieno[3,2-d]pyrimidin-2-yl]-1-hydroxy-vinyl]-4-fluoro-benzoni-
 trile;
 2-[2-[3-(2-Fluoro-phenyl)-4-oxo-3,4-dihydro-thieno[3,2-d]pyrimidin-2-yl]-1-hydroxy-vinyl]-4-methyl-benzoni-
 trile;
 2-[2-[3-(2-Chloro-phenyl)-4-oxo-3,4-dihydro-thieno[3,2-d]pyrimidin-2-yl]-1-hydroxy-vinyl]-benzonitrile; and,
 2-[2-[3-(2-Chloro-pyrid-3-yl)-4-oxo-3,4-dihydro-thieno[3,2-d]pyrimidin-2-yl]-1-hydroxy-vinyl]-benzonitrile;
 3-(2-chloro-phenyl)-6-fluoro-2-[2-hydroxy-2-(2-methyl-thiazol-4-yl)-vinyl]-3H-quinazolin-4-one;
 3-(2-chloro-phenyl)-6-fluoro-2-[2-hydroxy-2-(6-methyl-pyridin-2-yl)-vinyl]-3H-quinazolin-4-one;
 2-[2-[3-(2-chloro-phenyl)-6-fluoro-4-oxo-3,4-dihydro-quinazolin-2-yl]-1-hydroxy-vinyl]-6-methyl-nicotinoni-
 trile;
 2-[2-[3-(2-chloro-phenyl)-6-fluoro-4-oxo-3,4-dihydro-quinazolin-2-yl]-1-hydroxy-vinyl]-nicotinonitrile;
 2-[2-[3-(2-chloro-phenyl)-6-fluoro-4-oxo-3,4-dihydro-quinazolin-2-yl]-1-hydroxy-vinyl]-benzonitrile;
 2-[2-[3-(2-chloro-pyridin-3-yl)-6-fluoro-4-oxo-3,4-dihydro-quinazolin-2-yl]-1-hydroxy-vinyl]-6-methyl-nicoti-
 nonitrile;
 3-(2-chloro-phenyl)-6-fluoro-2-(2-hydroxy-2-pyridin-2-yl-vinyl)-3H-quinazolin-4-one;
 2-[2-[6-fluoro-3-(2-methyl-pyridin-3-yl)-4-oxo-3,4-dihydro-quinazolin-2-yl]-1-hydroxy-vinyl]-benzonitrile;
 2-[2-[3-(2-chloro-pyridin-3-yl)-6-fluoro-4-oxo-3,4-dihydro-quinazolin-2-yl]-1-hydroxy-vinyl]-benzonitrile;
 3-(2-chloro-phenyl)-6-fluoro-2-[2-(2-fluoro-phenyl)-2-hydroxy-ethyl]-3H-quinazolin-4-one;


(E) 3-(2-chloro-phenyl)-6-fluoro-2-[(pyridin-2-ylmethyl)-amino]-3H-quinazolin-4-one;

6-fluoro-3-(2-methyl-phenyl)-2-[(pyridin-2-ylmethyl)-amino]-3H-quinazolin-4-one;
 3-(2-chloro-phenyl)-6-fluoro-2-[(2-fluorophenyl-methyl)-amino]-3H-quinazolin-4-one;
 3-(2-chloro-phenyl)-2-[(2-cyanophenyl-methyl)-amino]-6-fluoro-3H-quinazolin-4-one;
 3-(2-chloro-phenyl)-2-[(6-diethylaminomethylpyridin-2-ylmethyl)-amino]-6-fluoro-3H-quinazolin-4-one;
 3-(2-chloro-phenyl)-6-fluoro-2-[(6-pyrrolidin-1-ylmethyl-pyridin-2-ylmethyl)-amino]-3H-quinazolin-4-one;
 3-(2-chloro-phenyl)-2-[(3-pyrrolidin-1-ylmethyl-phenylamino)-methyl]-3H-thieno[3,2-d]pyrimidin-4-one;
 3-(2-methyl-phenyl)-2-[(3-pyrrolidin-1-ylmethyl-phenylamino)-methyl]-3H-thieno[3,2-d]pyrimidin-4-one;
 3-(2-chloro-phenyl)-2-[(2-fluoro-phenylamino)-methyl]-3H-thieno[3,2-d]pyrimidin-4-one;
 3-(2-chloro-pyrid-3-yl)-2-[(3-pyrrolidin-1-ylmethyl-phenylamino)-methyl]-3H-thieno[3,2-d]pyrimidin-4-one;
 2-[(3-(2-chloro-pyrid-3-yl)-4-oxo-3,4-dihydro-thieno[3,2-d]pyrimidin-2-ylmethyl)-amino]-benzonitrile;
 3-(2-chloro-phenyl)-2-[(3-pyrrolidin-1-ylmethyl-phenylamino)-methyl]-3H-quinazolin-4-one;
 6-chloro-3-(2-chloro-phenyl)-2-[(3-pyrrolidin-1-ylmethyl-phenylamino)-methyl]-3H-quinazolin-4-one;
 6-chloro-3-(2-chloro-phenyl)-2-[(3-diethylaminomethyl-phenylamino)-methyl]-3H-quinazolin-4-one;
 6-chloro-3-(2-chloro-pyrid-3-yl)-2-[(3-diethylaminomethyl-phenylamino)-methyl]-3H-quinazolin-4-one;
 6-chloro-3-(2-trifluoromethyl-phenyl)-2-[(3-diethylaminomethyl-phenylamino)-methyl]-3H-quinazolin-4-one;
 2-[(3-(2-chloro-pyridin-3-yl)-4-oxo-3,4-dihydro-quinazolin-2-ylmethyl)-amino]-benzonitrile;


2-[[3-(2-methyl-pyridin-3-yl)-4-oxo-3,4-dihydro-quinazolin-2-ylmethyl]-amino]-benzonitrile;
 2-[[6-fluoro-3-(2-methyl-phenyl)-4-oxo-3,4-dihydro-quinazolin-2-ylmethyl]-amino]-nicotinonitrile;
 2-[[3-(2-chloro-phenyl)-4-oxo-3,4-dihydro-quinazolin-2-ylmethyl]-amino]-nicotinonitrile;
 2-[[3-(2-chloro-pyridin-3-yl)-6-fluoro-4-oxo-3,4-dihydro-quinazolin-2-ylmethyl]-amino]-benzonitrile;
 3-[[3-(2-chloro-phenyl)-6-fluoro-4-oxo-3,4-dihydro-quinazolin-2-ylmethyl]-amino]-benzonitrile;
 3-(2-chloro-phenyl)-2-[(3-diethylaminomethyl-phenylamino)-methyl]-6-fluoro-3H-quinazolin-4-one;
 3-(2-chloro-phenyl)-6-fluoro-2-(pyrimidin-2-ylaminomethyl)-3H-quinazolin-4-one;
 3-(2-chloro-pyridin-3-yl)-6-fluoro-2-(m-tolylamino-methyl)-3H-quinazolin-4-one;
 3-(2-chloro-pyridin-3-yl)-6-fluoro-2-[(6-methyl-pyridin-2-ylamino)-methyl]-3H-quinazolin-4-one;
 3-(2-chloro-phenyl)-6-fluoro-2-(pyridin-2-ylaminomethyl)-3H-quinazolin-4-one;
 3-(2-chloro-pyridin-3-yl)-6-fluoro-2-[(3-pyrrolidin-1-ylmethyl-phenylamino)-methyl]-3H-quinazolin-4-one;
 6-fluoro-3-(2-methyl-pyridin-3-yl)-2-[(3-pyrrolidin-1-ylmethyl-phenylamino)-methyl]-3H-quinazolin-4-one;
 3-(2-chloro-phenyl)-6-fluoro-2-[(2-fluoro-benzylamino)-methyl]-3H-quinazolin-4-one;
 N-[3-[(3-(2-chloro-phenyl)-6-fluoro-4-oxo-3,4-dihydro-quinazolin-2-ylmethyl)-amino]-phenyl]-acetamide;
 3-(2-chloro-phenyl)-6-fluoro-2-[(3-pyrrolidin-1-ylmethyl-phenylamino)-methyl]-3H-quinazolin-4-one;
 2-[[3-(2-chloro-phenyl)-6-fluoro-4-oxo-3,4-dihydro-quinazolin-2-ylmethyl]-amino]-nicotinonitrile;
 3-(2-chloro-pyridin-3-yl)-6-fluoro-2-[(2-fluoro-phenylamino)-methyl]-3H-quinazolin-4-one;
 3-(2-chloro-phenyl)-6-fluoro-2-[(2-fluoro-phenylamino)-methyl]-3H-quinazolin-4-one;
 3-(2-chloro-phenyl)-6-fluoro-2-[(6-methyl-pyridin-2-ylamino)-methyl]-3H-quinazolin-4-one; and,
 (F) an atropisomer of the formula



wherein either V, X, Y and Z are all carbon or one of them is nitrogen and the others are carbon;
 each of R¹, R², R³, R⁴ and R⁵ is selected, independently, from hydrogen, halogen, (C₁-C₆)alkyl, trifluoromethyl, cyano, (C₁-C₆)alkoxy, (C₁-C₆)alkylthio and C(=O)-O-(C₁-C₆)alkyl, with the proviso that: (a) R¹ can not be the same as R⁵, when each of V, X and Z is carbon; (b) at least one of R¹ and R⁵ must be other than hydrogen; and (c) when V, X, Y or Z is nitrogen, then R⁵, R⁴, R³ or R² respectively, is absent;
 ring A is a fused heteroaromatic ring, wherein said heteroaromatic ring is a 5 or 6 membered heteroaromatic ring, wherein said 6 membered heteroaromatic ring, taken together with the carbon atoms common to both rings of the bicyclic system, has the formula



and wherein said 5 membered heteroaromatic ring, taken together with the carbon atoms common to both rings of the bicyclic system, has the formula




wherein said ring positions "A", "B", "D" and "E" may be independently selected from carbon or nitrogen;


wherein said ring positions "F", "G" and "J" may be independently selected from carbon, nitrogen, oxygen or sulfur, with the proviso that: (a) if more than two of "F", "G" or "J" is a heteroatom then said 5 membered heteroaromatic ring is selected from the group consisting of (1,2,3)-triazole, (1,2,3)-thiadiazole, (1,2,5)-thiadiazole, and (1,2,5)-oxadiazole; and (b) if two of "F", "G" or "J" are heteroatoms, only one of said heteroatoms may be oxygen or sulfur;

wherein said fused heteroaromatic rings may optionally be independently substituted on any of the carbon or nitrogen atoms capable of forming an additional bond with a substituent selected from hydrogen, (C₁-C₆)alkyl, halogen, trifluoromethyl, amino-(CH₂)_n-, (C₁-C₆)alkylamino-(CH₂)_n-, di(C₁-C₆)alkyl-amino-(CH₂)_n-, (C₁-C₆)alkoxy, hydroxy(C₁-C₆)alkyl, (C₁-C₆)alkyl-O-(C₁-C₆)alkyl-, -CN, (C₁-C₆)alkyl-CO-O-(C₁-C₆)alkyl-, (C₁-C₆)alkyl-O-CO-O-(C₁-C₆)alkyl, (C₁-C₆)alkyl-CO-O-, hydroxy, -NO₂, R¹⁵-C(=O)-, R¹⁵-O-C(=O)-, di(C₁-C₆)alkyl-N-C(=O)-, (C₃-C₇)cycloalkyl, and R¹⁵-NH-C(=O)-, and phenyl optionally substituted with halo, (C₁-C₆)alkyl, -CN, or -CF₃;

[0006] R⁶ is phenyl of the formula Ph¹ or a five or six membered heterocycle, wherein said 6-membered heterocycle has the formula




wherein "N" is nitrogen; wherein said ring positions "K", "L" and "M" may be independently selected from carbon or nitrogen, with the proviso that only one of "K", "L" or "M" can be nitrogen; wherein said five membered heterocycle has the formula



wherein said ring positions "P," "Q" and "T" may be independently selected from carbon, nitrogen, oxygen or sulfur; with the proviso that only one of "P," "Q" or "T" can be oxygen or sulfur and at least one of "P," "Q" or "T" must be a heteroatom;

wherein said Ph¹ is a group of the formula



wherein each R¹⁵ is, independently, hydrogen or (C₁-C₆) alkyl;

each of R⁹, R¹⁰ and R¹¹ is selected, independently, from hydrogen, (C₁-C₆)alkyl optionally substituted with one to three halogen atoms, halo, CF₃, (C₁-C₆)alkoxy optionally substituted with one to three halogen atoms, (C₁-C₆)alkylthiol, R¹⁶O-(CH₂)_p-, (C₁-C₆)alkyl-NH-(CH₂)_p-, di(C₁-C₆)alkyl-N-(CH₂)_p-, (C₃-C₇)cycloalkyl-NH-(CH₂)_p-, H₂N-(C=O)-(CH₂)_p-, (C₁-C₆)alkyl-HN-(C=O)-(CH₂)_p-, di(C₁-C₆)alkyl-N-(C=O)-(CH₂)_p-, (C₃-C₇)cycloalkyl-NH-(C=O)-(CH₂)_p-, R¹⁶O-(C=O)-(CH₂)_p-, (C₁-C₆)alkyl-(O=C)-O-(C₁-C₆)alkyl-, (C₁-C₆)alkyl-O-(O=C)-O-(C₁-C₆)alkyl-, (C₁-C₆)alkyl-(O=C)-O-, (C₁-C₆)alkyl-(O=C)-NH-(CH₂)_p-, H(O=C)-NH-(CH₂)_p-, (C₁-C₆)alkyl-(O=C)-N-[(C₁-C₆)alkyl](CH₂)_p-, H(O=C)-N-[(C₁-C₆)alkyl](CH₂)_p-, hydroxy, H-C(=O)-(CH₂)_p-, (C₁-C₆)alkyl-C(=O)-, (C₁-C₆)alkyl-O-C(=O)-, R¹⁵-(CH₂)_p-O-C(=O)-, amino-(CH₂)_p-, hydroxy-(C₁-C₆)alkyl-, (C₁-C₆)alkyl-O-(C₁-C₆)alkyl- and cyano;

each of R⁷, R¹² and R¹³ is selected, independently, from hydrogen, (C₁-C₆)alkyl optionally substituted with one to three halogen atoms, halogen, CF₃, (C₁-C₆)alkoxy optionally substituted with one to three halogen atoms, (C₁-C₆)alkylthiol, R¹⁶O-(CH₂)_p-, (C₁-C₆)alkyl-NH-(CH₂)_p-, di(C₁-C₆)alkyl-N-(CH₂)_p-, (C₃-C₇)cycloalkyl-NH-(CH₂)_p-, H₂N-(C=O)-(CH₂)_p-, (C₁-C₆)alkyl-HN-(C=O)-(CH₂)_p-, di(C₁-C₆)alkyl-N-(C=O)-(CH₂)_p-, (C₃-C₇)cycloalkyl-NH-(C=O)-(CH₂)_p-, R¹⁶O-(C=O)-(CH₂)_p-, (C₁-C₆)alkyl-(O=C)-O-(C₁-C₆)alkyl-, (C₁-C₆)alkyl-O-(O=C)-O-(C₁-C₆)alkyl-, (C₁-C₆)alkyl-(O=C)-O-, (C₁-C₆)alkyl-(O=C)-NH-(CH₂)_p-, H(O=C)-NH-(CH₂)_p-, (C₁-C₆)alkyl-(O=C)-N-[(C₁-C₆)alkyl](CH₂)_p-, H(O=C)-N-[(C₁-C₆)alkyl](CH₂)_p-, hydroxy, H-C(=O)-(CH₂)_p-, (C₁-C₆)alkyl-C(=O)-, (C₁-C₆)alkyl-O-C(=O)-, R¹⁵-(CH₂)_p-O-C(=O)-, amino-(CH₂)_p-, hydroxy-(C₁-C₆)alkyl-, (C₁-C₆)alkyl-O-(C₁-C₆)alkyl-, -CHO and cyano;

each R¹⁴ is, independently, hydrogen or halogen;

each R¹⁶ is, independently, hydrogen, (C₁-C₆)alkyl, (C₁-C₆)alkyl-(C=O)-, (C₁-C₆)alkyl-O-(C=O)-, (C₁-C₆)alkyl-NH-(C=O)-, or di(C₁-C₆)alkyl-N-(C=O)-;

each is hydrogen, cyano, (C₁-C₆)alkyl, halogen, trifluoromethyl, -CHO or (C₁-C₆)alkoxy;

n is an integer from zero to 3;

p is an integer from zero to 3; and

wherein the dashed bond represented an optional double bond;

with the proviso that when R¹¹ is hydrogen, one of R¹³ and R¹⁴ is other than hydrogen.

[0007] In a specific embodiment of the above method, said dopamine agonist therapy is therapy comprising the administration of L-dopa or L-dopa in combination with an inhibitor of peripheral dopadecarboxylase such as carbidopa or benserazide.

[0008] In another specific embodiment of the above method, said compound is a compound of group (A) or a pharmaceutically acceptable salt thereof.

[0009] In another specific embodiment of the above method, said compound is a compound of group (B) or a pharmaceutically acceptable salt thereof.

[0010] This invention also relates to a method of treating dyskinesias associated with dopamine agonist therapy in a mammal, such as a human, which comprises administering to said mammal an AMPA receptor antagonizing effective amount of a compound within group (A), (B), (C), (D), (E), or (F), or a pharmaceutically acceptable salt of said compound, wherein groups (A), (B), (C), (D), (E), and (F) are as defined above.

compound, wherein groups (A), (B), (C), (D), (E), and (F) are as defined above.

[0011] This invention also relates to a method of treating dyskinesias associated with dopamine agonist therapy in a mammal, such as a human, which comprises administering to said mammal an AMPA receptor antagonizing effective amount of a compound selected from the group consisting of an AMPA receptor antagonist referred to in PCT international application publication number WO 97/19066; the compounds "NS-1201" or "NS-479" developed or marketed by Neurosearch (Denmark); the compound "LY-311446" (2-amino-3-(2-(3-(1H-tetrazol-5-yl)phenoxy)phenyl)propionic acid), "LY-300164" (7-acetyl-5-(4-aminophenyl)-8(R)-methyl-8,9-dihydro-7H-1,3-dioxolo(4,5-h)(2,3)benzodiazepine), "LY-293606", "LY-293558", or "GYKI-53655" of Eli Lilly (United States) or any AMPA antagonist referred to in 20th CINP (Melbourne), 1996, Abs S-40-1; the compound "NNC-07-0775" of Novo Nordisk (Denmark) or any AMPA antagonist referred to in PCT international publication number WO 96/15100; the compound "SYM-2206" (4-(aminophenyl)-1-methyl-6,7-(methylenedioxy)-N-butyl-1,2-dihydrophthalazine-2-carboxamide) of Symphony Pharmaceuticals (United States) or any AMPA antagonist referred to in Journal of Medicinal Chemistry, 1996, 39, 343; the compound "S-17625" (6,7-dichloro-2(1H)-oxoquinoline-3-phosphonic acid) of Servier (France) or any AMPA antagonist referred to in Journal of Medicinal Chemistry, 1996, 39, 197; 2-carboxy-1-methyl-7-trifluoromethylimidazo(1,2-a)quinoxalin-4(5H)-one or any AMPA antagonist referred to in PCT international publication numbers WO 95/21842, WO 96/08492, and WO 96/08493; 6-(4-pyridinyl)-1H-1,2,3-triazolo(4,5-a)pyrimidin-4(5H)-one or any AMPA antagonist referred to in Journal of Medicinal Chemistry 1995, 38, 587; any AMPA antagonist referred to in PCT international publication numbers WO 94/26747, WO 95/19346, WO 95/12594, WO 95/02601, WO 95/26342, WO 95/26349, WO 95/26350, WO 95/26351, WO 95/26352, WO 96/31511, and WO 95/02602; 2-amino-3-(3-hydroxy-5-(2-thienyl)isoxazol-4-yl)propionic acid or any AMPA antagonist referred to in PCT international publication number WO 95/12587; the compound "SYM-2250" of Symphony Pharmaceuticals (United States); the compound "S-18986" of Servier (France) or any AMPA antagonist referred to in 13th Int. Symp. Med. Chem. (Paris), 1994, Abs P29; the compound "NNC-07-9202" of Warner-Lambert (United States) or any AMPA antagonist referred to in 208th ACS (Washington, DC), 1994, Abs MEDI 170; the compound "IDRA-21" (7-chloro-3-methyl-3,4-dihydro-2H-1,2,4-benzothiadiazine-5,5-dioxide) or any AMPA antagonist referred to in Soc. Neurosci. Abs (Washington, DC), 1993, Abs 124.7 and 124.8; the compound "NS-409" of Warner-Lambert (United States) or any AMPA antagonist referred to in J. Med. Chem. 1995, 38, 3720 or PCT international publication numbers WO 96/08494 and WO 96/08495; the compound "NS-393" of Neurosearch (Denmark); the compounds "SYM-2101", "SYM-2007" and "SYM-2057" of Symphony Pharmaceuticals (United States); the compound "AMPAlex" (1-(1,3-benzodioxolo-5-ylcarbonyl)piperidine) of Cortex Pharmaceuticals (United States) or any AMPA antagonist referred to in Scrip, 1995, 2088/9, 14 and Scrip, 1996, 2187, 21 or in PCT international publication number WO 96/38414; the compounds "LY-293558", "LY-215490", and decahydro-6-(2-(1H-tetrazol-5-yl)ethyl)-3-isoxolinol-1-carboxylic acid (CAS registry no. 154652-83-2) or any AMPA antagonist referred to in J. Med. Chem., 1993, 36, 2046; the compound "YM-90K" (1,4-dihydro-6-(1H-imidazol-1-yl)-7-nitro-2,3-quinoxalinedione monohydrochloride (CAS registry no. 154164-30-4 or any AMPA antagonist referred to in Scrip, 1994, 1972, 14 or PCT international publication number WO 96/10023; the compound "aloracetam" (N-(2-(3-formyl-2,5-dimethyl-1H-pyrrol-1-yl)ethyl)-acetamide)(CAS registry no. 119610-26-3) or any AMPA antagonist referred to in European Patent 287988; the compound "NS-257" of Warner-Lambert; the compound "NNC-07-9202" of Novo Nordisk (Denmark) or any AMPA antagonist referred to in European Patent 283959 and Science, 1988, 241, 701; and the compound "aniracetam" of Roche (Switzerland) or 1-(4-methoxybenzyl)-2-pyrrolidinone (CAS registry no. 72432-10-1) or any AMPA antagonist referred to in European Patent 5143.

[0012] The term "treating", as used herein, unless otherwise indicated, means reversing, alleviating, inhibiting the progress of, or preventing the disorder or condition to which such term applies, or one or more symptoms of such disorder or condition. The term "treatment", as used herein, refers to the act of treating, as "treating" is defined immediately above.



[0013] The term "dyskinesia(s)", as used herein, unless otherwise indicated, means any abnormal or uncontrollable

[0014] The term or phrase "dopamine agonist therapy", as used herein, unless otherwise indicated, means any therapy that increases dopamine receptor stimulation, including, but not limited to, therapies that directly stimulate dopamine receptors (such as bromocriptine) and therapies that increase the levels of dopamine (such as L-dopa or drugs which inhibit dopamine metabolism). Dopamine agonist therapies include, but are not limited to, therapies comprising the administration of one or more of the following agents: L-dopa, L-dopa in combination with an L-dopa decarboxylase inhibitor such as carbidopa or benserazide, bromocriptine, dihydroergocryptine, etisulergine, AF-14, alaptide, pergolide, piribedil, dopamine D1 receptor agonists such as A-68939, A-77636, dihydrexine, and SKF-38393;

dopamine D2 receptor agonists such as carbergoline, lisuride, N-0434, naxagolide, PD-118440, pramipexole, quinpirole and ropinirole; dopamine/ β -adrenergic receptor agonists such as DPDMS and dopexamine; dopamine/5-HT uptake inhibitor/5-HT-1A agonists such as roxindole; dopamine/opiate receptor agonists such as NIH-10494; α 2-adrenergic antagonist/dopamine agonists such as terguride; α 2-adrenergic antagonist/dopamine D2 agonists such as ergolines and talipexole; dopamine uptake inhibitors such as GBR-12909, GBR-13069, GYKI-52895, and NS-2141; monoamine oxidase-B inhibitors such as selegiline, N-(2-butyl)-N-methylpropargylamine, N-methyl-N-(2-pentyl)propargylamine, AGN-1133, ergot derivatives, lazabemide, LU-53439, MD-280040 and mofegiline; and COMT inhibitors such as CGP-28014, entacapone and tolcapone. Dopamine agonist therapy, as referred to in the present invention, is used in the treatment of a central nervous system disorder such as, but not limited to, Parkinson's disease.

[0015] The term or phrase "dyskinesia associated with dopamine agonist therapy", as used herein, unless otherwise indicated, means any dyskinesia which accompanies, or follows in the course of, dopamine agonist therapy, or which is caused by, related to, or exacerbated by dopamine agonist therapy, wherein dyskinesia and dopamine agonist therapy are as defined above.

[0016] In the compounds of groups (A) and (B), referred to above, the designation "(S)" appearing at the beginning of each compound refers to the configuration of each compound as an atropisomer. The compounds of group (F) are also atropisomers, and the compounds of groups (C), (D) and (E) include atropisomers. Atropisomers are conformational isomers that occur when rotation about a single bond in the molecule is prevented or greatly slowed as a result of steric interactions with other parts of the molecule and the substituents at both ends of the single bond are unsymmetrical. A detailed account of atropisomers can be found in Jerry March, *Advanced Organic Chemistry*, 101-102 (4th ed. 1992) and in Oki, *Top. Stereochem.*, 14, 1-81 (1983). Each compound within groups (A), (B) and (F) has the same (S) configuration as an atropisomer. This configuration is described in United States provisional patent applications numbers 60/038905 (filed February 28, 1997) and 60/038540 (filed February 28, 1997), both of which are referred to above. This configuration may be illustrated with respect to the first compound listed in group (A) which is (S)-3-(2-chlorophenyl)-2-[2-(5-diethylaminomethyl-2-fluorophenyl)-vinyl]-6-fluoro-3H-quinazolin-4-one. Below, both atropisomeric configurations are illustrated.



[0017] In the above structures, the bold lines indicate that the bolded atoms of the 2-chlorophenyl group are sterically restricted so as to exist above the plane of the quinazolinone ring. This steric restriction is due to a rotational energy barrier preventing free rotation about the single bond connecting the nitrogen at position 3 of the quinazolinone ring to the 2-chlorophenyl group. The above (S) configuration is also illustrated in formula I of group (F). The other compounds of groups (A), (B) and (F) are all atropisomers having an (S) configuration analogous to the structure labeled "(S) Configuration" illustrated above. The compounds of groups (C), (D), and (E) also exist, and may be isolated as, atropisomers having (S) and (R) configurations corresponding to the (S) and (R) configurations illustrated above.

[0018] In addition to the atropisomerism referred to above, the compounds of groups (A), (B), (C), (D), (E), and (F) may have chiral centers and therefore may exist in different enantiomeric and diastereomeric forms. This invention relates to all optical isomers and all stereoisomers of compounds of groups (A), (B), (C), (D), (E), and (F), and mixtures thereof, and to all methods of treatment defined above that contain or employ them, respectively.

[0019] The method of the present invention also relates to the use of pharmaceutically acceptable acid addition salts of the compounds of groups (A), (B), (C), (D), (E), and (F). The acids which are used to prepare the pharmaceutically acceptable acid addition salts of the aforementioned base compounds of this invention are those which form non-toxic acid addition salts, i.e., salts containing pharmacologically acceptable anions, such as the hydrochloride, hydrobromide, hydroiodide, nitrate, sulfate, bisulfate, phosphate, acid phosphate, acetate, lactate, citrate, acid citrate, tartrate,

bitartrate, succinate, maleate, fumarate, gluconate, saccharate, benzoate, methanesulfonate, ethanesulfonate, benzenesulfonate, p-toluenesulfonate and pamoate [i.e., 1,1'-methylene-bis-(2-hydroxy-3-naphthoate)] salts

[0020] The invention also relates to base addition salts of the compounds of groups (A), (B), (C), (D), (E), and (F). The chemical bases that may be used as reagents to prepare pharmaceutically acceptable base salts of those compounds of groups (A), (B), (C), (D), (E), and (F) that are acidic in nature are those that form non-toxic base salts with such compounds. Such non-toxic base salts include, but are not limited to those derived from such pharmacologically acceptable cations such as alkali metal cations (e.g., potassium and sodium) and alkaline earth metal cations (e.g., calcium and magnesium), ammonium or water-soluble amine addition salts such as N-methylglucamine (meglumine), and the lower alkanolammonium and other base salts of pharmaceutically acceptable organic amines.

Detailed Description Of The Invention

[0021] The compounds of groups (A), (B), (C), (D), (E), and (F) are readily prepared. The compounds of group (A) can be prepared and separated as atropisomers according to one or more methods referred to in PCT/IB98/00150 referred to above. The compounds of group (B) can be prepared and separated as atropisomers according to one or more methods referred to in PCT/IB98/00151, referred to above. The compounds of group (C) can be prepared according to one or more methods referred to in PCT international application number PCT/IB97/00134 (publication no. WO 9743276), referred to above. The compounds of group (D) can be prepared according to one or more methods referred to in EP98304522.0, referred to above. The compounds of group (E) can be prepared according to one or more methods referred to in EP98304319.0, referred to above. The compounds of group (F) can be prepared according to one or more methods referred to in United States patent application (provisional no. 60/057990) entitled "Novel Atropisomers Of 2,3-Disubstituted-(5,6)-Heteroaryl-fused-Pyrimidin-4-ones" filed July 23, 1998 with Bertrand L. Chenard and Willard M. Welch named as inventors, referred to above.

[0022] The compounds of groups (A), (B), (C), (D), (E), and (F), referred to above, which are basic in nature are capable of forming a wide variety of different salts with various inorganic and organic acids. Although such salts must be pharmaceutically acceptable for administration to animals, it is often desirable in practice to initially isolate a compound of group (A), (B), (C), (D), (E), or (F) from the reaction mixture as a pharmaceutically unacceptable salt and then simply convert the latter back to the free base compound by treatment with an alkaline reagent, and subsequently convert the free base to a pharmaceutically acceptable acid addition salt. The acid addition salts of the base compounds of the method of this invention are readily prepared by treating the base compound with a substantially equivalent amount of the chosen mineral or organic acid in an aqueous solvent medium or in a suitable organic solvent such as methanol or ethanol. Upon careful evaporation of the solvent, the desired solid salt is obtained.

[0023] The acids which are used to prepare the pharmaceutically acceptable acid addition salts, i.e., salts containing compounds of groups (A), (B), (C), (D), (E), and (F) are those which form non-toxic acid addition salts, i.e., salts containing pharmacologically acceptable anions, such as hydrochloride, hydrobromide, hydroiodide, nitrate, sulfate or bisulfate, phosphate or acid phosphate, acetate, lactate, citrate or acid citrate, tartrate or bitartrate, succinate, maleate, fumarate, gluconate, saccharate, benzoate, methanesulfonate and pamoate [i.e., 1,1'-methylene-bis-(2-hydroxy-3-naphthoate)] salts.

[0024] Those compounds of groups (A), (B), (C), (D), (E), and (F) which are acidic in nature are capable of forming base salts with various pharmacologically acceptable cations. Examples of such salts include the alkali metal or alkaline-earth metal salts and particular, the sodium and potassium salts. These salts are all prepared by conventional techniques. The chemical bases which are used as reagents to prepare the pharmaceutically acceptable base salts of this invention are those which form non-toxic base salts with the herein described acidic compounds of groups (A), (B), (C), (D), (E), and (F). These non-toxic base salts include those derived from such pharmacologically acceptable cations as sodium, potassium, calcium and magnesium, etc. These salts can easily be prepared by treating the responding acidic compounds with an aqueous solution containing the desired pharmacologically acceptable cations, and then evaporating the resulting solution to dryness, preferably under reduced pressure. Alternatively, they may also be prepared by mixing lower alkanolic solutions of the acidic compounds and the desired alkali metal alkoxide together, and then evaporating the resulting solution to dryness in the same manner as before. In either case, stoichiometric quantities of reagents are preferably employed in order to ensure completeness of reaction of maximum product of yields of the desired final product.

[0025] The *in vitro* and *in vivo* activity of the compounds of groups (A), (B), (C), (D), (E), and (F) for AMPA receptor antagonism can be determined by methods available to one of ordinary skill in the art. One method for determining the activity of the compounds of groups (A), (B), (C), (D), (E), and (F) is by blockage of AMPA receptor activation-induced $^{45}\text{Ca}^{2+}$ uptake into neurons. A specific method for determining blockage of AMPA receptor activation-induced $^{45}\text{Ca}^{2+}$ uptake into neurons is described below.

Neuronal primary cultures

[0026] Primary cultures of rat cerebellar granule neurons are prepared as described by Parks, T.N., Artman, L.D., Alasti, N., and Nemeth, E.F., Modulation Of N-Methyl-D-Aspartate Receptor-Mediated Increases In Cytosolic Calcium In Cultured Rat Cerebellar Granule Cells, Brain Res. 552, 13-22 (1991). According to this method, cerebella are removed from 8 day old CD rats, minced into 1 mm pieces and incubated for 15 minutes at 37°C in calcium-magnesium free Tyrode's solution containing 0.1 % trypsin. The tissue is then triturated using a fine bore Pasteur pipette. The cell suspension is plated onto poly-D-lysine coated 96-well tissue culture plates at 10^5 cells per well. Medium consists of Minimal Essential Medium (MEM), with Earle's salts, 10% heat inactivated Fetal Bovine Serum, 2 mM L-glutamine, 21 mM glucose, Penicillin-Streptomycin (100 units per ml) and 25 mM KCl. After 24 hours, the medium is replaced with fresh medium containing 10 μ M cytosine arabinoside to inhibit cell division. Cultures are used 6 to 8 days later.

AMPA receptor activation-induced $^{45}\text{Ca}^{2+}$ uptake

[0027] The effects of drugs on AMPA receptor activation-induced $^{45}\text{Ca}^{2+}$ uptake can be examined in rat cerebellar granule cell cultures prepared as described above. Cultures in 96 well plates are preincubated for approximately 3 hours in serum free medium and then for 10 minutes in a Mg^{2+} -free balanced salt solution (in mM: 120 NaCl, 5 KCl, 0.33 NaH_2PO_4 , 1.8 CaCl_2 , 22.0 glucose and 10.0 HEPES at pH 7.4) containing 0.5 mM DTT, 10 μ M glycine and drugs at 2X final concentration. The reaction is started by rapid addition of an equal volume of the balanced salt solution containing 100 μ M of the AMPA receptor agonist kainic acid and $^{45}\text{Ca}^{2+}$ (final specific activity 250 Ci/mmol). After 10 minutes at 25°C, the reaction is stopped by aspirating the $^{45}\text{Ca}^{2+}$ -containing solution and washing the cells 5X in an ice cold balanced salt solution containing no added calcium and 0.5 mM EDTA. Cells are then lysed by overnight incubation in 0.1 % Triton-X100 and radioactivity in the lysate is then determined. All compounds of groups (A), (B), (C), (D), (E), and (F), referred to above, at concentrations of 0.5 μ M or less inhibited the AMPA receptor activation-induced $^{45}\text{Ca}^{2+}$ uptake by 50% or more.

[0028] The following procedure may be used to assess the efficacy of the compounds of groups (A), (B), (C), (D), (E), and (F) in the treatment of dyskinesias associated with dopamine agonist therapy in the treatment of Parkinson's disease. Aged, female rhesus monkeys are rendered Parkinsonian as follows. Each monkey is first infused with 0.4 mg/kg MPTP (1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine) via the right internal carotid artery. After being evaluated behaviorally for 3 to 6 weeks and being judged to have stable unilateral deficits, the animals receive a second MPTP injection via the left internal carotid artery. Monkeys lesioned according to this protocol have been shown to have stable, bilateral deficits that are responsive to L-dopa and apomorphine. Once the monkeys are Parkinsonian, dyskinesias are induced over a period of approximately 3 to 6 weeks by treating the monkeys twice daily with subcutaneous injections of PHNO ((+)-4-propyl-9-hydroxynaphthoxazine) (a dopamine agonist). Dyskinesias are assessed 30 minutes after PHNO injection and every 30 minutes for the next 120 minutes (5 measurements) taking into account the following: type of dyskinesia (chorea, dystonia); intensity (0 = absent; 1 = mild; 2 = moderate; 3 = severe); and topography (arm, leg, trunk, generalized). The overall score (0 - 3) is averaged over the 5 measurements. Scoring is performed blindly from coded videotapes. A compound of group (A), (B), (C), (D), (E), or (F) is then administered together with the dopamine agonist at dosages ranging from 0.05 mg/kg to 1 mg/kg.

[0029] Pharmaceutical compositions for use in the method of the present invention may be prepared according to methods familiar to those skilled in the art. For example, pharmaceutical compositions containing a compound of group (A), (B), (C), (D), (E), or (F), or a pharmaceutically acceptable salt thereof (hereinafter the "active compound(s)") may be formulated in a conventional manner using one or more pharmaceutically acceptable carriers. Thus, the active compounds may be formulated for oral, buccal, intranasal, parenteral (e.g., intravenous, intramuscular or subcutaneous), transdermal (e.g., patch, ointment, cream or iontophoresis), or rectal administration or in a form suitable for administration by inhalation or insufflation.

[0030] For oral administration, the pharmaceutical compositions may take the form of, for example, tablets or capsules prepared by conventional means with pharmaceutically acceptable excipients such as binding agents (e.g., pregelatinised maize starch, polyvinylpyrrolidone or hydroxypropyl methylcellulose); fillers (e.g., lactose, microcrystalline cellulose or calcium phosphate); lubricants (e.g., magnesium stearate, talc or silica); disintegrants (e.g., potato starch or sodium starch glycolate); or wetting agents (e.g., sodium lauryl sulphate). The tablets may be coated by methods well known in the art. Liquid preparations for oral administration may take the form of, for example, solutions, syrups or suspensions, or they may be presented as a dry product for constitution with water or other suitable vehicle before use. Such liquid preparations may be prepared by conventional means with pharmaceutically acceptable additives such as suspending agents (e.g., sorbitol syrup, methyl cellulose or hydrogenated edible fats); emulsifying agents (e.g., lecithin or acacia); non-aqueous vehicles (e.g., almond oil, oily esters or ethyl alcohol); and preservatives (e.g., methyl or propyl p-hydroxybenzoates or sorbic acid).

[0031] For buccal administration, the pharmaceutical composition may take the form of tablets or lozenges formulated

in conventional manner.

[0032] The active compounds may be formulated for parenteral administration by injection, including using conventional catheterization techniques or infusion. Formulations for injection may be presented in unit dosage form, e.g., in ampoules or in multi-dose containers, with an added preservative. The compositions may take such forms as suspensions, solutions or emulsions in oily or aqueous vehicles, and may contain formulating agents such as suspending, stabilizing and/or dispersing agents. Alternatively, the active ingredient may be in powder form for reconstitution with a suitable vehicle, e.g., sterile pyrogen-free water, before use.

[0033] The active compounds may also be formulated in rectal compositions such as suppositories or retention enemas, e.g., containing conventional suppository bases such as cocoa butter or other glycerides.

[0034] For intranasal administration or administration by inhalation, the active compounds are conveniently delivered in the form of a solution or suspension from a pump spray container that is squeezed or pumped by the patient or as an aerosol spray presentation from a pressurized container or a nebulizer, with the use of a suitable propellant, e.g., dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, carbon dioxide or other suitable gas. In the case of a pressurized aerosol, the dosage unit may be determined by providing a valve to deliver a metered amount. The pressurized container or nebulizer may contain a solution or suspension of the active compound. Capsules and cartridges (made, for example, from gelatin) for use in an inhaler or insufflator may be formulated containing a powder mix of an active compound and a suitable powder base such as lactose or starch.

[0035] A proposed dose of the active compounds for use in the method of the present invention for oral, parenteral or buccal administration to the average adult human requiring treatment is 0.01 to 100 mg/kg of the active ingredient per unit dose which could be administered, for example, 1 to 4 times per day.

[0036] Aerosol formulations for use in the method of the present invention in the treatment of an average adult human are preferably arranged so that each metered dose or "puff" of aerosol contains 20 µg to 1000 µg of the active compound. The overall daily dose with an aerosol will be within the range 100 µg to 10 mg. Administration may be several times daily, for example 2, 3, 4 or 8 times, giving for example, 1, 2 or 3 doses each time.

[0037] For transdermal administration the composition may take the form of patches, creams, ointments or iontophoresis formulated in conventional manner such as described in United States Patents 5,004,610 and 5,364,630, issued April 2, 1991 and November 15, 1994 respectively.

Claims

1. The use of a compound selected from groups (A), (B), (C), (D), (E), or (F) or a pharmaceutically acceptable salt thereof, in the manufacture of a medicament for treating dyskinesia associated with dopamine agonist therapy, wherein groups (A), (B), (C), (D), (E), and (F) are defined as follows:

(A) (S)-3-(2-chloro-phenyl)-2-[2-(5-diethylaminomethyl-2-fluoro-phenyl)-vinyl]-6-fluoro-3H-quinazolin-4-one;

(S)-3-(2-chloro-phenyl)-2-[2-(6-diethylaminomethyl-pyridin-2-yl)-vinyl]-6-fluoro-3H-quinazolin-4-one;

(S)-3-(2-chloro-phenyl)-2-[2-(4-diethylaminomethyl-pyridin-2-yl)-vinyl]-6-fluoro-3H-quinazolin-4-one;

(S)-3-(2-chloro-phenyl)-2-[2-(6-ethylaminomethyl-pyridin-2-yl)-vinyl]-6-fluoro-3H-quinazolin-4-one;

(S)-3-(2-bromo-phenyl)-2-[2-(6-diethylaminomethyl-pyridin-2-yl)-vinyl]-6-fluoro-3H-quinazolin-4-one;

(S)-3-(2-chloro-phenyl)-6-fluoro-2-[2-(6-methoxymethyl-pyridin-2-yl)-vinyl]-3H-quinazolin-4-one;

(S)-3-(2-chloro-phenyl)-6-fluoro-2-[2-(4-methyl-pyrimidine-2-yl)-vinyl]-3H-quinazolin-4-one;

(S)-3-(2-chloro-phenyl)-6-fluoro-2-[2-[6-(isopropylamino-methyl)-pyridin-2-yl]-ethyl]-3H-quinazolin-

4-one;

(S)-6-fluoro-2-[2-(2-methyl-thiazol-4-yl)-vinyl]-3-(2-methyl-phenyl)-3H-quinazolin-4-one;

(S)-3-(2-chloro-phenyl)-6-fluoro-2-[2-(2-methyl-thiazol-4-yl)-vinyl]-3H-quinazolin-4-one;

(S)-2-[2-(2-dimethylaminomethyl-thiazol-4-yl)-vinyl]-6-fluoro-3-(2-fluoro-phenyl)-3H-quinazolin-4-one;

(S)-3-(2-bromo-phenyl)-6-fluoro-2-[2-(2-methyl-thiazol-4-yl)-vinyl]-3H-quinazolin-4-one;

(S)-3-(2-chloro-phenyl)-2-[2-(2-methyl-thiazol-4-yl)-vinyl]-3H-quinazolin-4-one;

(S)-3-(2-chloro-phenyl)-6-fluoro-2-(2-pyridin-2-yl-vinyl)-3H-quinazolin-4-one;

(S)-3-(2-bromo-phenyl)-2-(2-pyridin-2-yl-vinyl)-3H-quinazolin-4-one;

(S)-6-chloro-2-(2-pyridin-2-yl-vinyl)-3-o-tolyl-3H-quinazolin-4-one;

(S)-3-(2-chloro-phenyl)-2-[2-(6-methyl-pyridin-2-yl)-vinyl]-3H-quinazolin-4-one;

(S)-6-chloro-2-[2-(6-methyl-pyridin-2-yl)-vinyl]-3-o-tolyl-3H-quinazolin-4-one;

(S)-3-(2-chloro-phenyl)-6-fluoro-2-(2-pyridin-2-yl-ethyl)-3H-quinazolin-4-one;

(S)-6-[2-[3-(2-chloro-phenyl)-6-fluoro-4-oxo-3,4-dihydro-quinazolin-2-yl]-vinyl]-pyridine-2-carbaldehyde;

(S)-3-(2-chloro-phenyl)-6-fluoro-2-[2-(6-methylaminomethyl-pyridin-2-yl)-vinyl]-3H-quinazolin-4-one;

(S)-N-(6-{2-[3-(2-chloro-phenyl)-6-fluoro-4-oxo-3,4-dihydro-quinazolin-2-yl]-vinyl}-pyridin-2-ylmethyl)-N-methyl-acetamide;

(S)-6-{2-[3-(2-chloro-phenyl)-6-fluoro-4-oxo-3,4-dihydro-quinazolin-2-yl]-vinyl}-pyridine-2-carbonitrile;

(S)-3-(2-fluoro-phenyl)-2-(2-pyridin-2-yl-vinyl)-3H-quinazolin-4-one;

(S)-3-(2-bromo-phenyl)-6-fluoro-2-(2-pyridin-2-yl-vinyl)-3H-quinazolin-4-one;

(S)-3-(4-bromo-2-chloro-phenyl)-6-fluoro-2-(2-pyridin-2-yl-vinyl)-3H-quinazolin-4-one;

(S)-3-(2-chloro-phenyl)-2-[2-(6-diethylaminomethyl-pyridin-2-yl)-vinyl]-3H-quinazolin-4-one;

(S)-N-(6-{2-[3-(2-chloro-phenyl)-6-fluoro-4-oxo-3,4-dihydro-quinazolin-2-yl]-vinyl}-pyridin-2-ylmethyl)-N-ethyl-acetamide;

(S)-3-(2-chloro-phenyl)-6-fluoro-2-[2-(6-fluoromethyl-pyridin-2-yl)-vinyl]-3H-quinazolin-4-one;

(S)-3-(2-chloro-phenyl)-6-fluoro-2-[2-(6-pyrrolidin-1-ylmethyl-pyridin-2-yl)-ethyl]-3H-quinazolin-4-one;

(S)-3-(2-chloro-phenyl)-2-[2-(6-{ethyl-(2-hydroxy-ethyl)-amino}-methyl)-pyridin-2-yl]-vinyl]-6-fluoro-3H-quinazolin-4-one;

(S)-3-(2-chloro-phenyl)-6-fluoro-2-[2-(6-(isopropylamino-methyl)-pyridin-2-yl)-vinyl]-3H-quinazolin-4-one;

(S)-3-(2-chloro-phenyl)-6-fluoro-2-[2-(6-(2-methyl-piperidin-1-ylmethyl)-pyridin-2-yl)-vinyl]-3H-quinazolin-4-one;

(S)-3-(2-chloro-phenyl)-2-[2-(6-ethoxymethyl-pyridin-2-yl)-vinyl]-6-fluoro-3H-quinazolin-4-one;

(S)-3-(2-chloro-phenyl)-2-[2-(6-(2,5-dihydro-pyrrol-1-ylmethyl)-pyridin-2-yl)-vinyl]-6-fluoro-3H-quinazolin-4-one;

(S)-3-(2-chloro-phenyl)-6-fluoro-2-[2-(6-(4-methyl-piperidin-1-ylmethyl)-pyridin-2-yl)-vinyl]-3H-quinazolin-4-one;

(S)-6-bromo-2-[2-(6-methyl-pyridin-2-yl)-vinyl]-3-o-tolyl-3H-quinazolin-4-one;

(S)-6-bromo-2-(2-pyridin-2-yl-vinyl)-3-o-tolyl-3H-quinazolin-4-one;

(S)-6-fluoro-3-(2-fluoro-phenyl)-2-(2-pyridin-2-yl-vinyl)-3H-quinazolin-4-one;

(S)-3-(2-chloro-phenyl)-6-methyl-2-(2-pyridin-2-yl-vinyl)-3H-quinazolin-4-one;

(S)-6-fluoro-3-(2-fluoro-phenyl)-2-[2-(6-dimethylaminomethyl-pyridin-2-yl)-vinyl]-6-fluoro-3H-quinazolin-4-one;

(S)-6-fluoro-3-(2-fluoro-phenyl)-2-[2-(6-methyl-pyridin-2-yl)-vinyl]-3H-quinazolin-4-one;

(S)-3-(2-chloro-phenyl)-2-[2-(6-{(2-dimethylamino-ethyl)-methyl-amino}-methyl)-pyridin-2-yl)-vinyl]-6-fluoro-3H-quinazolin-4-one;

(S)-3-(2-chloro-phenyl)-6-fluoro-2-[2-(6-hydroxymethyl-pyridin-2-yl)-vinyl]-3H-quinazolin-4-one;

(S)-acetic acid 6-{2-[3-(2-chloro-phenyl)-6-fluoro-4-oxo-3,4-dihydro-quinazolin-2-yl]-vinyl}-pyridin-2-yl methyl ester;

(S)-6-{2-[3-(2-bromo-phenyl)-6-fluoro-4-oxo-3,4-dihydro-quinazolin-2-yl]-vinyl}-pyridine-2-carbaldehyde;

(S)-3-(2-bromo-phenyl)-2-[2-(6-diethylaminomethyl-pyridin-2-yl)-vinyl]-3H-quinazolin-4-one;

(S)-acetic acid 6-{2-[3-(2-bromo-phenyl)-6-fluoro-4-oxo-3,4-dihydro-quinazolin-2-yl]-vinyl}-pyridin-2-yl methyl ester;

(S)-diethylamino-acetic acid 6-{2-[3-(2-chloro-phenyl)-6-fluoro-4-oxo-3,4-dihydro-quinazolin-2-yl]-vinyl}-pyridin-2-ylmethyl ester;

(S)-3-(2-chloro-phenyl)-2-[2-(6-difluoromethyl-pyridin-2-yl)-vinyl]-6-fluoro-3H-quinazolin-4-one;

(S)-3-(2-chloro-phenyl)-6-fluoro-2-[2-(6-methoxy-pyridin-2-yl)-vinyl]-3H-quinazolin-4-one;

(S)-2-[2-[3-(2-chloro-phenyl)-6-fluoro-4-oxo-3,4-dihydro-quinazolin-2-yl]-vinyl]-6-methyl-nicotinonitrile;

(S)-2-[2-[3-(2-chloro-phenyl)-6-fluoro-4-oxo-3,4-dihydro-quinazolin-2-yl]-ethyl]-6-methyl-nicotinonitrile;

(S)-3-(2-chloro-phenyl)-6-fluoro-2-(2-pyrimidine-2-yl-ethyl)-3H-quinazolin-4-one;

(S)-3-(2-chloro-phenyl)-2-[2-(4,6-dimethyl-pyrimidine-2-yl)-vinyl]-6-fluoro-3H-quinazolin-4-one;

(S)-2-[2-[3-(2-chloro-phenyl)-6-fluoro-4-oxo-3,4-dihydro-quinazolin-2-yl]-vinyl]-nicotinonitrile;

(S)-3-(2-chloro-phenyl)-6-fluoro-2-(2-[6-(3-methyl-butylamino)-methyl]-pyridin-2-yl)-ethyl)-3H-quinazolin-4-one;

(S)-2-[2-[3-(2-chloro-phenyl)-6-fluoro-4-oxo-3,4-dihydro-quinazolin-2-yl]-ethyl]-nicotinonitrile;

(S)-2-[2-(6-chloro-4-oxo-3-o-tolyl-3,4-dihydro-quinazolin-2-yl)-vinyl]-benzonitrile;

(S)-2-[2-[3-(2-chloro-phenyl)-6-fluoro-4-oxo-3,4-dihydro-quinazolin-2-yl]-vinyl]-4-methyl-benzonitrile;

(S)-3-(2-bromo-phenyl)-6-fluoro-2-[2-(6-hydroxymethyl-pyridin-2-yl)-vinyl]-3H-quinazolin-4-one; and

(S)-3-(2-chloro-phenyl)-6-fluoro-2-[2-(6-pyrrolidin-1-ylmethyl-pyridin-2-yl)-vinyl]-3H-quinazolin-4-one;

(B) (S)-6-fluoro-2-[2-(2-fluoro-phenyl)-vinyl]-3-(2-methyl-pyridin-3-yl)-3H-quinazolin-4-one;

(S)-2-[2-[6-fluoro-3-(2-methyl-pyridin-3-yl)-4-oxo-3,4-dihydro-quinazolin-2-yl]-vinyl]-benzonitrile;

(S)-2-[2-[6-fluoro-3-(2-methylpyridin-3-yl)-4-oxo-3,4-dihydroquinazolin-2-yl]-vinyl]-benzonitrile;

(S)-2-{2-[3-(2-chloro-pyridin-3-yl)-6-fluoro-4-oxo-3,4-dihydroquinazolin-2-yl]-vinyl}-benzonitrile;
 (S)-2-{2-[6-fluoro-3-(2-methyl-pyridin-3-yl)-4-oxo-3,4-dihydro-quinazolin-2-yl]-vinyl}-4-methyl-benzoni-
 trile;
 (S)-2-{2-[3-(2-methyl-pyridin-3-yl)-4-oxo-3,4-dihydro-quinazolin-2-yl]vinyl}-benzonitrile;
 (S)-6-fluoro-3-(2-methyl-pyridin-3-yl)-2-[2-(thiazol-2-yl)-vinyl]-3H-quinazolin-4-one;
 (S)-6-fluoro-3-(2-methyl-pyridin-3-yl)-2-[2-(2-methyl-thiazol-4-yl)-vinyl]-3H-quinazolin-4-one;
 (S)-6-fluoro-3-(2-methyl-pyridin-3-yl)-2-[2-(4-methyl-thiazol-2-yl)-vinyl]-3H-quinazolin-4-one;
 (S)-2-[2-(5-diethylaminomethyl-2-fluoro-phenyl)-vinyl]-6-fluoro-3-(2-methyl-pyridin-3-yl)-3H-quinazolin-
 4-one;
 (S)-6-fluoro-2-[2-(2-fluoro-5-pyrrolidin-1-ylmethyl-phenyl)-vinyl]-3-(2-methyl-pyridin-3-yl)-3H-quinazolin-
 4-one;
 (S)-3-(2-chloro-pyridin-3-yl)-2-[2-(2-fluoro-phenyl)-vinyl]-3H-quinazolin-4-one;
 (S)-3-(2-chloro-pyridin-3-yl)-6-fluoro-2-[2-(6-methyl-phenyl-2-yl)-vinyl]-3H-quinazolin-4-one;
 (S)-3-(2-chloro-pyridin-3-yl)-6-fluoro-2-[2-(fluoro-phenyl)-vinyl]-3H-quinazolin-4-one;
 (S)-6-chloro-2-[2-(2-fluoro-phenyl)-vinyl]-3-(2-methyl-pyridin-3-yl)-3H-quinazolin-4-one;
 (S)-6-chloro-2-[2-(2-fluoro-phenyl)-vinyl]-3-(3-methyl-1-oxy-pyridin-4-yl)-3H-quinazolin-4-one;
 (S)-3-[2-(3-(2-chloro-pyridin-3-yl)-6-fluoro-4-oxo-3,4-dihydro-quinazolin-2-yl)-vinyl]-benzaldehyde;
 (S)-3-[2-[3-(2-chloro-pyridin-3-yl)-4-oxo-3,4-dihydro-quinazolin-2-yl]-vinyl]-benzaldehyde;
 (S)-3-(2-chloro-pyridin-3-yl)-6-fluoro-2-[2-(3-hydroxymethyl-phenyl)-vinyl]-3H-quinazolin-4-one;
 (S)-3-(2-chloro-pyridin-3-yl)-2-[2-[3(1,4-dioxo-8-aza-spiro[4.5]dec-8-ylmethyl)-phenyl]-vinyl]-6-fluoro-
 3H-quinazolin-4-one;
 (S)-3-(2-chloro-pyridin-3-yl)-6-fluoro-2-[2-[3-(4-pyrrolidin-1-yl-piperidin-1-ylmethyl)-phenyl]-vinyl]-3H-
 quinazolin-4-one;
 (S)-2-{2-[3-(2-chloro-pyridin-3-yl)-6-fluoro-4-oxo-3,4-dihydro-quinazolin-2-yl]-vinyl}-benzonitrile;
 (S)-2-{2-[3-(2-chloro-pyridin-3-yl)-4-oxo-3,4-dihydro-quinazolin-2-yl]-vinyl}-benzonitrile;
 (S)-2-[2-(2-fluoro-phenyl)-vinyl]-3-(2-methyl-pyridin-3-yl)-3H-quinazolin-4-one;
 (S)-3-(2-chloro-pyridin-3-yl)-6-fluoro-2-[2-hydroxy-phenyl]-vinyl]-3H-quinazolin-4-one;
 (S)-6-fluoro-3-(2-methyl-pyridin-3-yl)-2-[2-(2-methyl-thiazol-4-yl)-ethyl]-3H-quinazolin-4-one;
 (S)-6-fluoro-3-(2-chloro-pyridin-3-yl)-2-[2-(2-dimethylamino-methylthiazol-4-yl)-vinyl]-3H-quinazolin-
 4-one;
 (S)-2-[2-(5-Diethylaminomethyl-2-fluoro-phenyl)-vinyl]-6-fluoro-3-(4-methyl-pyridin-3-yl)-3H-quinazolin-
 4-one;
 (S)-4-Diethylaminomethyl-2-[2-[6-fluoro-3-(4-methyl-pyridin-3-yl)-4-oxo-3,4-dihydro-quinazolin-2-yl]-vi-
 nyl]-benzonitrile;
 (S)-2-[2-(5-Diethylaminomethyl-2-fluoro-phenyl)-vinyl]-6-fluoro-3-(3-methyl-pyrazin-2-yl)-3H-
 quinazolin-4-one;
 (S)-6-fluoro-3-(2-methyl-pyridin-3-yl)-2-[2-(2-dimethylamino-methylthiazol-4-yl)-vinyl]-3H-quinazolin-
 4-one;
 (S)-6-fluoro-3-(2-methyl-pyridin-3-yl)-2-[2-(2-methyl-oxazol-4-yl)-vinyl]-3H-quinazolin-4-one;
 (S)-6-fluoro-3-(2-chloro-pyridin-3-yl)-2-[2-(thiazol-2-yl)-vinyl]-3H-quinazolin-4-one;
 (S)-6-fluoro-3-(4-methyl-pyridin-3-yl)-2-[2-(4-methyl-thiazol-2-yl)-vinyl]-3H-quinazolin-4-one;
 (S)-3-(2-chloro-pyridin-3-yl)-6-fluoro-2-[2-(2-hydroxy-phenyl)-vinyl]-3H-quinazolin-4-one; and
 (S)-6-fluoro-2-[2-(2-fluoro-5-pyrrolidin-1-ylmethyl-phenyl)-ethyl]-3-(2-methyl-pyridin-3-yl)-3H-
 quinazolin-4-one;

(C) 3-(2-chloro-phenyl)-6-fluoro-2-(2-pyridin-2-yl-vinyl)-3H-quinazolin-4-one;

3-(2-bromo-phenyl)-2-(2-pyridin-2-yl-vinyl)-3H-quinazolin-4-one;
 6-chloro-2-(2-pyridin-2-yl-vinyl)-3-o-tolyl-3H-quinazolin-4-one;
 3-(2-chloro-phenyl)-2-[2-(6-methyl-pyridin-2-yl)-vinyl]-3H-quinazolin-4-one;
 6-chloro-2-[2-(6-methyl-pyridin-2-yl)-vinyl]-3-o-tolyl-3H-quinazolin-4-one;
 3-(2-chloro-phenyl)-6-fluoro-2-(2-pyridin-2-yl-ethyl)-3H-quinazolin-4-one;
 6-[2-[3-(2-chloro-phenyl)-6-fluoro-4-oxo-3,4-dihydro-quinazolin-2-yl]-vinyl]-pyridine-2-carbaldehyde;
 3-(2-chloro-phenyl)-6-fluoro-2-[2-(6-methylaminomethyl-pyridin-2-yl)-vinyl]-3H-quinazolin-4-one;
 N-(6-[2-[3-(2-chloro-phenyl)-6-fluoro-4-oxo-3,4-dihydro-quinazolin-2-yl]-vinyl]-pyridin-2-ylmethyl)-N-me-
 thyl-acetamide;
 3-(2-chloro-phenyl)-2-[2-(4-diethylaminomethyl-pyridin-2-yl)-vinyl]-6-fluoro-3H-quinazolin-4-one;
 6-[2-[3-(2-chloro-phenyl)-6-fluoro-4-oxo-3,4-dihydro-quinazolin-2-yl]-vinyl]-pyridine-2-carbonitrile;

3-(2-fluoro-phenyl)-2-(2-pyridin-2-yl-vinyl)-3H-quinazolin-4-one;
 3-(2-bromo-phenyl)-6-fluoro-2-(2-pyridin-2-yl-vinyl)-3H-quinazolin-4-one;
 3-(4-bromo-2-chloro-phenyl)-6-fluoro-2-(2-pyridin-2-yl-vinyl)-3H-quinazolin-4-one;
 3-(2-chloro-phenyl)-2-[2-(6-diethylaminomethyl-pyridin-2-yl)-vinyl]-3H-quinazolin-4-one;
 N-(6-{2-[3-(2-chloro-phenyl)-6-fluoro-4-oxo-3,4-dihydro-quinazolin-2-yl]-vinyl}-pyridin-2-ylmethyl)-N-ethyl-acetamide;
 3-(2-chloro-phenyl)-6-fluoro-2-[2-(6-fluoromethyl-pyridin-2-yl)-vinyl]-3H-quinazolin-4-one;
 3-(2-chloro-phenyl)-6-fluoro-2-[2-(6-pyrrolidin-1-ylmethyl-pyridin-2-yl)-ethyl]-3H-quinazolin-4-one;
 3-(2-chloro-phenyl)-2-[2-(6-{ethyl-(2-hydroxy-ethyl)-amino}-methyl)-pyridin-2-yl]-vinyl]-6-fluoro-3H-quinazolin-4-one;
 3-(2-chloro-phenyl)-6-fluoro-2-[2-[6-(isopropylamino-methyl)-pyridin-2-yl]-vinyl]-3H-quinazolin-4-one;
 3-(2-chloro-phenyl)-6-fluoro-2-[2-[6-(2-methyl-piperidin-1-ylmethyl)-pyridin-2-yl]-vinyl]-3H-quinazolin-4-one;
 3-(2-chloro-phenyl)-2-[2-(6-ethylaminomethyl-pyridin-2-yl)-vinyl]-6-fluoro-3H-quinazolin-4-one;
 3-(2-chloro-phenyl)-2-[2-(6-ethoxymethyl-pyridin-2-yl)-vinyl]-6-fluoro-3H-quinazolin-4-one;
 3-(2-chloro-phenyl)-2-[2-[6-(2,5-dihydro-pyrrol-1-ylmethyl)-pyridin-2-yl]-vinyl]-6-fluoro-3H-quinazolin-4-one;
 3-(2-chloro-phenyl)-6-fluoro-2-[2-[6-(4-methyl-piperidin-1-ylmethyl)-pyridin-2-yl]-vinyl]-3H-quinazolin-4-one;
 6-bromo-2-[2-(6-methyl-pyridin-2-yl)-vinyl]-3-o-tolyl-3H-quinazolin-4-one;
 6-bromo-2-(2-pyridin-2-yl-vinyl)-3-o-tolyl-3H-quinazolin-4-one;
 6-fluoro-3-(2-fluoro-phenyl)-2-(2-pyridin-2-yl-vinyl)-3H-quinazolin-4-one;
 1-benzyl-5-(2-methyl-[1,3]dioxolan-2-yl)-2-oxo-2,3-dihydro-1H-indole-3-carboxylic acid (3-phenylcarbamoyl-phenyl)-amide;
 3-(2-chloro-phenyl)-6-methyl-2-(2-pyridin-2-yl-vinyl)-3H-quinazolin-4-one;
 3-(2-chloro-phenyl)-2-[2-(6-dimethylaminomethyl-pyridin-2-yl)-vinyl]-6-fluoro-3H-quinazolin-4-one;
 6-fluoro-3-(2-fluoro-phenyl)-2-[2-(6-methyl-pyridin-2-yl)-vinyl]-3H-quinazolin-4-one;
 3-(2-chloro-phenyl)-2-[2-(6-{[(2-dimethylamino-ethyl)-methyl-amino]-methyl}-pyridin-2-yl)-vinyl]-6-fluoro-3H-quinazolin-4-one;
 3-(2-chloro-phenyl)-6-fluoro-2-[2-(6-hydroxymethyl-pyridin-2-yl)-vinyl]-3H-quinazolin-4-one;
 acetic acid 6-{2-[3-(2-chloro-phenyl)-6-fluoro-4-oxo-3,4-dihydro-quinazolin-2-yl]-vinyl}-pyridin-2-ylmethyl ester;
 6-{2-[3-(2-bromo-phenyl)-6-fluoro-4-oxo-3,4-dihydro-quinazolin-2-yl]-vinyl}-pyridine-2-carbaldehyde;
 3-(2-bromo-phenyl)-2-[2-(6-diethylaminomethyl-pyridin-2-yl)-vinyl]-6-fluoro-3H-quinazolin-4-one;
 3-(2-bromo-phenyl)-2-[2-(6-diethylaminomethyl-pyridin-2-yl)-vinyl]-3H-quinazolin-4-one;
 acetic acid 6-{2-[3-(2-bromo-phenyl)-6-fluoro-4-oxo-3,4-dihydro-quinazolin-2-yl]-vinyl}-pyridin-2-ylmethyl ester;
 3-(2-chloro-phenyl)-6-fluoro-2-[2-(6-methoxymethyl-pyridin-2-yl)-vinyl]-3H-quinazolin-4-one;
 diethylamino-acetic acid 6-{2-[3-(2-chloro-phenyl)-6-fluoro-4-oxo-3,4-dihydroquinazolin-2-yl]-vinyl}-pyridin-2-ylmethyl ester;
 6-fluoro-3-(2-methyl-pyridin-3-yl)-2-[2-(2-methyl-thiazol-4-yl)-vinyl]-3H-quinazolin-4-one;
 3-(2-bromo-phenyl)-6-fluoro-2-[2-(6-hydroxymethyl-pyridin-2-yl)-vinyl]-3H-quinazolin-4-one; and
 3-(2-chloro-phenyl)-6-fluoro-2-[2-(6-pyrrolidin-1-ylmethyl-pyridin-2-yl)-vinyl]-3H-quinazolin-4-one;

(D) 3-(2-chloro-phenyl)-2-[2-(6-diethylaminomethyl)-pyridin-2-yl]-2-hydroxy-vinyl]-6-fluoro-3H-quinazolin-4-one

6-Chloro-3-(2-chloro-phenyl)-2-[2-hydroxy-2-(6-methyl-pyridin-2-yl)-vinyl]-3H-quinazolin-4-one;
 2-[2-[3-(2-Chloro-phenyl)-4-oxo-3,4-dihydro-quinazolin-2-yl]-1-hydroxy-vinyl]-nicotinonitrile;
 2-[2-[3-(2-Chloro-pyrid-3-yl)-6-fluoro-4-oxo-3,4-dihydro-quinazolin-2-yl]-1-hydroxy-vinyl]-nicotinonitrile;
 2-[2-[6-Chloro-3-(2-methyl-phenyl)-4-oxo-3,4-dihydro-quinazolin-2-yl]-1-hydroxy-vinyl]-nicotinonitrile;
 3-(2-Chloro-phenyl)-2-[2-(3-diethylaminomethyl-phenyl)-2-hydroxy-ethyl]-6-fluoro-3H-quinazolin-4-one;
 3-(2-Chloro-phenyl)-6-fluoro-2-[2-(3-pyrrolidin-1-ylmethyl-phenyl)-2-hydroxy-ethyl]-3H-quinazolin-4-one;
 3-(2-Chloro-pyrid-3-yl)-2-[2-(3-diethylaminomethyl-phenyl)-2-hydroxy-ethyl]-6-fluoro-3H-quinazolin-4-one;
 2-[2-(3-Diethylaminomethyl-phenyl)-2-hydroxy-ethyl]-6-fluoro-3-(2-fluoro-phenyl)-3H-quinazolin-4-one;
 2-[2-(3-Diethylaminomethyl-phenyl)-2-hydroxy-ethyl]-3-(2-fluoro-phenyl)-3H-quinazolin-4-one;
 3-(2-chloro-phenyl)-2-[2-(6-diethylaminomethyl)-pyridin-2-yl]-2-hydroxy-vinyl]-6-fluoro-3H-quinazolin-4-one;


2-{2-[3-(2-Chloro-pyrid-3-yl)-6-fluoro-4-oxo-3,4-dihydro-quinazolin-2-yl]-1-hydroxy-vinyl}-6-methyl-nico-
 tinonitrile;
 2-{2-[3-(2-Chloro-phenyl)-4-oxo-3,4-dihydro-quinazolin-2-yl]-1-hydroxy-vinyl}-6-methyl-nicotinonitrile;
 2-{2-[6-Chloro-3-(2-chloro-phenyl)-4-oxo-3,4-dihydro-quinazolin-2-yl]-1-hydroxy-vinyl}-6-methyl-nicoti-
 5 nonitrile;
 2-{2-[3-(2-Chloro-phenyl)-6-fluoro-4-oxo-3,4-dihydro-quinazolin-2-yl]-1-hydroxy-vinyl}-6-fluoro-nicoti-
 nonitrile;
 2-{2-[3-(2-Chloro-phenyl)-6-fluoro-4-oxo-3,4-dihydro-quinazolin-2-yl]-1-hydroxy-vinyl}-4-fluoro-benzoni-
 10 trile;
 2-{2-[3-(2-Chloro-phenyl)-6-fluoro-4-oxo-3,4-dihydro-quinazolin-2-yl]-1-hydroxy-vinyl}-4-methyl-ben-
 zonitrile;
 2-{2-[3-(2-Chloro-phenyl)-4-oxo-3,4-dihydro-thieno[3,2-d]pyrimidin-2-yl]-1-hydroxy-vinyl}-6-methyl-nico-
 tinonitrile;
 2-{2-[3-(2-methyl-phenyl)-4-oxo-3,4-dihydro-thieno[3,2-d]pyrimidin-2-yl]-1-hydroxy-vinyl}-6-methyl-nico-
 15 tinonitrile;
 2-{2-[3-(2-Chloro-pyrid-3yl)-4-oxo-3,4-dihydro-thieno[3,2-d]pyrimidin-2-yl]-1-hydroxyvinyl}-4-methyl-
 benzonitrile;
 2-{2-[3-(2-Chloro-phenyl)-4-oxo-3,4-dihydro-thieno[3,2-d]pyrimidin-2-yl]-1-hydroxy-vinyl}-4-fluoro-ben-
 zonitrile;
 20 2-{2-[3-(2-Fluoro-phenyl)-4-oxo-3,4-dihydro-thieno[3,2-d]pyrimidin-2-yl]-1-hydroxyvinyl}-4-methyl-ben-
 zonitrile;
 2-{2-[3-(2-Chloro-phenyl)-4-oxo-3,4-dihydro-thieno[3,2-d]pyrimidin-2-yl]-1-hydroxyvinyl}-benzonitrile;
 and,
 2-{2-[3-(2-Chloro-pyrid-3yl)-4-oxo-3,4-dihydro-thieno[3,2-d]pyrimidin-2-yl]-1-hydroxyvinyl}-benzonitrile;
 25 3-(2-chloro-phenyl)-6-fluoro-2-[2-hydroxy-2-(2-methyl-thiazol-4-yl)-vinyl]-3H-quinazolin-4-one;
 3-(2-chloro-phenyl)-6-fluoro-2-[2-hydroxy-2-(6-methyl-pyridin-2-yl)-vinyl]-3H-quinazolin-4-one;
 2-{2-[3-(2-chloro-phenyl)-6-fluoro-4-oxo-3,4-dihydro-quinazolin-2-yl]-1-hydroxy-vinyl}-6-methyl-nicoti-
 nonitrile;
 2-{2-[3-(2-chloro-phenyl)-6-fluoro-4-oxo-3,4-dihydro-quinazolin-2-yl]-1-hydroxy-vinyl}-nicotinonitrile;
 30 2-{2-[3-(2-chloro-phenyl)-6-fluoro-4-oxo-3,4-dihydro-quinazolin-2-yl]-1-hydroxy-vinyl}-benzonitrile;
 2-{2-[3-(2-chloro-pyridin-3-yl)-6-fluoro-4-oxo-3,4-dihydro-quinazolin-2-yl]-1-hydroxy-vinyl}-6-methyl-
 nicotinonitrile;
 3-(2-chloro-phenyl)-6-fluoro-2-(2-hydroxy-2-pyridin-2-yl-vinyl)-3H-quinazolin-4-one;
 2-{2-[6-fluoro-3-(2-methyl-pyridin-3-yl)-4-oxo-3,4-dihydro-quinazolin-2-yl]-1-hydroxy-vinyl}-benzonitrile;
 35 2-{2-[3-(2-chloro-pyridin-3-yl)-6-fluoro-4-oxo-3,4-dihydro-quinazolin-2-yl]-1-hydroxy-vinyl}-benzonitrile;
 3-(2-chloro-phenyl)-6-fluoro-2-[2-(2-fluoro-phenyl)-2-hydroxy-ethyl]-3H-quinazolin-4-one;

(E) 3-(2-chloro-phenyl)-6-fluoro-2-[(pyridin-2-ylmethyl)-amino]-3H-quinazolin-4-one;


6-fluoro-3-(2-methyl-phenyl)-2-[(pyridin-2-ylmethyl)-amino]-3H-quinazolin-4-one;
 3-(2-chloro-phenyl)-6-fluoro-2-[(2-fluorophenyl-methyl)-amino]-3H-quinazolin-4-one;
 3-(2-chloro-phenyl)-2-[(2-cyanophenyl-methyl)-amino]-6-fluoro-3H-quinazolin-4-one;
 3-(2-chloro-phenyl)-2-[(6-diethylaminomethylpyridin-2-ylmethyl)-amino]-6-fluoro-3H-quinazolin-4-one;
 3-(2-chloro-phenyl)-6-fluoro-2-[(6-pyrrolidin-1-ylmethyl-pyridin-2-ylmethyl)-amino]-3H-quinazolin-4-one;
 45 3-(2-chloro-phenyl)-2-[(3-pyrrolidin-1-ylmethyl-phenylamino)-methyl]-3H-thieno[3,2-d]pyrimidin-4-one;
 3-(2-methyl-phenyl)-2-[(3-pyrrolidin-1-ylmethyl-phenylamino)-methyl]-3H-thieno[3,2-d]pyrimidin-4-one;
 3-(2-chloro-phenyl)-2-[(2-fluoro-phenylamino)-methyl]-3H-thieno[3,2-d]pyrimidin-4-one;
 3-(2-chloro-pyrid-3-yl)-2-[(3-pyrrolidin-1-ylmethyl-phenylamino)-methyl]-3H-thieno[3,2-d]pyrimidin-
 4-one;
 50 2-{[3-(2-chloro-pyrid-3-yl)-4-oxo-3,4-dihydro-thieno[3,2-d]pyrimidin-2-ylmethyl]-amino}-benzonitrile;
 3-(2-chloro-phenyl)-2-[(3-pyrrolidin-1-ylmethyl-phenylamino)-methyl]-3H-quinazolin-4-one;
 6-chloro-3-(2-chloro-phenyl)-2-[(3-pyrrolidin-1-ylmethyl-phenylamino)-methyl]-3H-quinazolin-4-one;
 6-chloro-3-(2-chloro-phenyl)-2-[(3-diethylaminomethyl-phenylamino)-methyl]-3H-quinazolin-4-one;
 6-chloro-3-(2-chloro-pyrid-3-yl)-2-[(3-diethylaminomethyl-phenylamino)-methyl]-3H-quinazolin-4-one;
 55 6-chloro-3-(2-trifluoromethyl-phenyl)-2-[(3-diethylaminomethyl-phenylamino)-methyl]-3H-quinazolin-
 4-one;
 2-{[3-(2-chloro-pyridin-3-yl)-4-oxo-3,4-dihydro-quinazolin-2-ylmethyl]-amino}-benzonitrile;
 2-{[3-(2-methyl-pyridin-3-yl)-4-oxo-3,4-dihydro-quinazolin-2-ylmethyl]-amino}-benzonitrile;

2-[[6-fluoro-3-(2-methyl-phenyl)-4-oxo-3,4-dihydro-quinazolin-2-ylmethyl]-amino]-nicotinonitrile;
 2-[[3-(2-chloro-phenyl)-4-oxo-3,4-dihydro-quinazolin-2-ylmethyl]-amino]-nicotinonitrile;
 2-[[3-(2-chloro-pyridin-3-yl)-6-fluoro-4-oxo-3,4-dihydro-quinazolin-2-ylmethyl]-amino]-benzonitrile;
 3-[[3-(2-chloro-phenyl)-6-fluoro-4-oxo-3,4-dihydro-quinazolin-2-ylmethyl]-amino]-benzonitrile;
 3-(2-chloro-phenyl)-2-[(3-diethylaminomethyl-phenylamino)-methyl]-6-fluoro-3H-quinazolin-4-one;
 3-(2-chloro-phenyl)-6-fluoro-2-(pyrimidin-2-ylaminomethyl)-3H-quinazolin-4-one;
 3-(2-chloro-pyridin-3-yl)-6-fluoro-2-(m-tolylamino-methyl)-3H-quinazolin-4-one;
 3-(2-chloro-pyridin-3-yl)-6-fluoro-2-[(6-methyl-pyridin-2-ylamino)-methyl]-3H-quinazolin-4-one;
 3-(2-chloro-phenyl)-6-fluoro-2-(pyridin-2-ylaminomethyl)-3H-quinazolin-4-one;
 3-(2-chloro-pyridin-3-yl)-6-fluoro-2-[(3-pyrrolidin-1-ylmethyl-phenylamino)-methyl]-3H-quinazolin-4-one;
 6-fluoro-3-(2-methyl-pyridin-3-yl)-2-[(3-pyrrolidin-1-ylmethyl-phenylamino)-methyl]-3H-quinazolin-4-one;
 3-(2-chloro-phenyl)-6-fluoro-2-[(2-fluoro-benzylamino)-methyl]-3H-quinazolin-4-one;
 N-{3-[[3-(2-chloro-phenyl)-6-fluoro-4-oxo-3,4-dihydro-quinazolin-2-ylmethyl]-amino]-phenyl}-acetamide;
 3-(2-chloro-phenyl)-6-fluoro-2-[(3-pyrrolidin-1-ylmethyl-phenylamino)-methyl]-3H-quinazolin-4-one;
 2-[[3-(2-chloro-phenyl)-6-fluoro-4-oxo-3,4-dihydro-quinazolin-2-ylmethyl]-amino]-nicotinonitrile;
 3-(2-chloro-pyridin-3-yl)-6-fluoro-2-[(2-fluoro-phenylamino)-methyl]-3H-quinazolin-4-one;
 3-(2-chloro-phenyl)-6-fluoro-2-[(2-fluoro-phenylamino)-methyl]-3H-quinazolin-4-one;
 3-(2-chloro-phenyl)-6-fluoro-2-[(6-methyl-pyridin-2-ylamino)-methyl]-3H-quinazolin-4-one; and,


(F) an atropisomer of the formula



wherein either V, X, Y and Z are all carbon or one of them is nitrogen and the others are carbon;
 each of R¹, R², R³, R⁴ and R⁵ is selected, independently, from hydrogen, halogen, (C₁-C₆)alkyl, trifluoromethyl, cyano, (C₁-C₆)alkoxy, (C₁-C₆)alkylthio and C(=O)-O-(C₁-C₆)alkyl, with the proviso that: (a) R¹ can not be the same as R⁵, when each of V, X and Z is carbon; (b) at least one of R¹ and R⁵ must be other than hydrogen; and (c) when V, X, Y or Z is nitrogen, then R⁵, R⁴, R³ or R² respectively, is absent;
 ring A is a fused heteroaromatic ring, wherein said heteroaromatic ring is a 5 or 6 membered heteroaromatic ring, wherein said 6 membered heteroaromatic ring, taken together with the carbon atoms common to both rings of the bicyclic system, has the formula



and wherein said 5 membered heteroaromatic ring, taken together with the carbon atoms common to both rings of the bicyclic system, has the formula




wherein said ring positions "A", "B", "D" and "E" may be independently selected from carbon or nitrogen;


wherein said ring positions "F", "G" and "J" may be independently selected from carbon, nitrogen, oxygen or sulfur, with the proviso that: (a) if more than two of "F", "G" or "J" is a heteroatom then said 5 membered heteroaromatic ring is selected from the group consisting of (1,2,3)-thiadiazole, (1,2,5)-thiadiazole, and (1,2,5)-oxadiazole; and (b) if two of "F", "G" or "J" are heteroatoms, only one of said heteroatoms may be oxygen or sulfur;

wherein said fused heteroaromatic rings may optionally be independently substituted on any of the carbon or nitrogen atoms capable of forming an additional bond with a substituent selected from hydrogen, (C₁-C₆)alkyl, halogen, trifluoromethyl, amino-(CH₂)_n, (C₁-C₆)alkylamino-(CH₂)_n, di(C₁-C₆)alkyl-amino-(CH₂)_n, (C₁-C₆)alkoxy, hydroxy(C₁-C₆)alkyl, (C₁-C₆)alkyl-O-(C₁-C₆)alkyl-, -CN, (C₁-C₆)alkyl-CO-O-(C₁-C₆)alkyl-, (C₁-C₆)alkyl-O-CO-O-(C₁-C₆)alkyl, (C₁-C₆)alkyl-CO-O-, hydroxy, -NO₂, R¹⁵-C(=O)-, R¹⁵-O-C(=O)-, di(C₁-C₆)alkyl-N-C(=O)-, (C₃-C₇)cycloalkyl, and R¹⁵-NH-C(=O)-, and phenyl optionally substituted with halo, (C₁-C₆)alkyl, -CN, or -CF₃;

R⁶ is phenyl of the formula Ph¹ or a five or six membered heterocycle, wherein said 6-membered heterocycle has the formula




wherein "N" is nitrogen; wherein said ring positions "K", "L" and "M" may be independently selected from carbon or nitrogen, with the proviso that only one of "K", "L" or "M" can be nitrogen; wherein said five membered heterocycle has the formula



wherein said ring positions "P," "Q" and "T" may be independently selected from carbon, nitrogen, oxygen or sulfur; with the proviso that only one of "P," "Q" or "T" can be oxygen or sulfur and at least one of "P," "Q" or "T" must be a heteroatom;

wherein said Ph¹ is a group of the formula



wherein each R¹⁵ is, independently, hydrogen or (C₁-C₆) alkyl;

each of R⁹, R¹⁰ and R¹¹ is selected, independently, from hydrogen, (C₁-C₆)alkyl optionally substituted with one to three halogen atoms, halo, CF₃, (C₁-C₆)alkoxy optionally substituted with one to three halogen atoms, (C₁-C₆)alkylthiol, R¹⁶O-(CH₂)_p-, (C₁-C₆)alkyl-NH-(CH₂)_p-, di(C₁-C₆)alkyl-N-(CH₂)_p-, (C₃-C₇)cycloalkyl-NH-(CH₂)_p-, H₂N-(C=O)-(CH₂)_p-, (C₁-C₆)alkyl-HN-(C=O)-(CH₂)_p-, di(C₁-C₆)alkyl-N-(C=O)-(CH₂)_p-, (C₃-C₇)cycloalkyl-NH-(C=O)-(CH₂)_p-, R¹⁶O-(C=O)-(CH₂)_p-, (C₁-C₆)alkyl-(O=C)-O-(C₁-C₆)alkyl-, (C₁-C₆)alkyl-O-(O=C)-O-(C₁-C₆)alkyl-, (C₁-C₆)alkyl-(O=C)-O-, (C₁-C₆)alkyl-(O=C)-NH-(CH₂)_p-, H(O=C)-NH-(CH₂)_p-, (C₁-C₆)alkyl-(O=C)-N-[(C₁-C₆)alkyl](CH₂)_p-, H(O=C)-N-[(C₁-C₆)alkyl](CH₂)_p-, hydroxy, H-C(=O)-(CH₂)_p-, (C₁-C₆)alkyl-C(=O)-, (C₁-C₆)alkyl-O-C(=O)-, R¹⁵-(CH₂)_p-O-C(=O)-, amino-(CH₂)_p-, hydroxy-(C₁-C₆)alkyl-, (C₁-C₆)alkyl-O-(C₁-C₆)alkyl- and cyano;

each of R⁷, R¹² and R¹³ is selected, independently, from hydrogen, (C₁-C₆)alkyl optionally substituted with one to three halogen atoms, halogen, CF₃, (C₁-C₆)alkoxy optionally substituted with one to three halogen atoms, (C₁-C₆)alkylthiol, R¹⁶O-(CH₂)_p-, (C₁-C₆)alkyl-NH-(CH₂)_p-, di(C₁-C₆)alkyl-N-(CH₂)_p-, (C₃-C₇)cycloalkyl-NH-(CH₂)_p-, H₂N-(C=O)-(CH₂)_p-, (C₁-C₆)alkyl-HN-(C=O)-(CH₂)_p-, di(C₁-C₆)alkyl-N-(C=O)-(CH₂)_p-, (C₃-C₇)cycloalkyl-NH-(C=O)-(CH₂)_p-, R¹⁶O-(C=O)-(CH₂)_p-, (C₁-C₆)alkyl-(O=C)-O-(C₁-C₆)alkyl-, (C₁-C₆)alkyl-O-(O=C)-O-(C₁-C₆)alkyl-, (C₁-C₆)alkyl-(O=C)-O-, (C₁-C₆)alkyl-(O=C)-NH-(CH₂)_p-, H(O=C)-NH-(CH₂)_p-, (C₁-C₆)alkyl-(O=C)-N-[(C₁-C₆)alkyl](CH₂)_p-, H(O=C)-N-[(C₁-C₆)alkyl](CH₂)_p-, hydroxy, H-C(=O)-(CH₂)_p-, (C₁-C₆)alkyl-C(=O)-, (C₁-C₆)alkyl-O-C(=O)-, R¹⁵-(CH₂)_p-O-C(=O)-, amino-(CH₂)_p-, hydroxy-(C₁-C₆)alkyl-, (C₁-C₆)alkyl-O-(C₁-C₆)alkyl-, -CHO and cyano;

each R¹⁴ is, independently, hydrogen or halogen;

each R¹⁶ is, independently, hydrogen, (C₁-C₆)alkyl, (C₁-C₆)alkyl-C(=O)-, (C₁-C₆)alkyl-O-C(=O)-, (C₁-C₆)alkyl-NH-C(=O)-, or di(C₁-C₆)alkyl-N-(C=O)-;

each is hydrogen, cyano, (C₁-C₆)alkyl, halogen, trifluoromethyl, -CHO or (C₁-C₆)alkoxy;

n is an integer from zero to 3;


p is an integer from zero to 3; and

wherein the dashed bond represented an optional double bond;

with the proviso that when R¹¹ is hydrogen, one of R¹³ and R¹⁴ is other than hydrogen.

2. The use as claimed in claim 1 wherein said dopamine agonist therapy is therapy comprising the administration of L-dopa or L-dopa in combination with an inhibitor of peripheral dopadecarboxylase

3. The use as claimed in claim 2 wherein said inhibitor of peripheral dopadecarboxylase is carbidopa or benserazide.
4. The use as claimed in any one of the preceding claims wherein said compound is a compound of group (A) or a pharmaceutically acceptable salt thereof.
5. The use as claimed in any one of claims 1 to 4 wherein said compound is a compound of group (B) or a pharmaceutically acceptable salt thereof.
6. The use as claimed in any one of the preceding claims wherein an AMPA receptor antagonizing effective amount of a compound within groups (A) to (F) is used.
7. The use as claimed in any one of the preceding claims wherein said compounds are used for the treatment of dyskinesias associated with dopamine agonist therapy in the treatment of Parkinson's disease.



(19)

Europäisches Patentamt
European Patent Office
Office européen des brevets



(11)

EP 0 900 568 A3

(12)

EUROPEAN PATENT APPLICATION

(88) Date of publication A3:
02.05.2001 Bulletin 2001/18

(51) Int Cl.7: **A61K 31/505**
// (A61K31/505, 31:195, 31:15)

(43) Date of publication A2:
10.03.1999 Bulletin 1999/10

(21) Application number: 98307181.2

(22) Date of filing: 04.09.1998

(84) Designated Contracting States:
AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE
Designated Extension States:
AL LT LV MK RO SI

- Menniti, Frank Samuel
Mystic, Connecticut 06355 (US)
- Welch, Willard KcKowan, Jr.
Mystic, Connecticut 6355 (US)

(30) Priority: 05.09.1997 US 58098 P

(71) Applicant: **Pfizer Products Inc.**
Groton, CT 06340-5146 (US)

(74) Representative: **McMunn, Watson Palmer et al**
Pfizer Limited
Patents Department
Ramsgate Road
Sandwich, Kent CT13 9NJ (GB)

(72) Inventors:
• **Chenard, Bertrand Leo**
Waterford, Connecticut 06385 (US)

(54) **AMPA antagonists for the treatment of dyskinesias associated with dopamine agonist therapy**

(57) The invention relates to a method of treating dyskinesias associated with dopamine agonist therapy in a mammal which comprises administering to said mammal a compound, as defined herein, which is an

antagonist of the AMPA receptor. Dopamine agonist therapy, as referred to in the present invention, is generally used in the treatment of a central nervous system disorder such as Parkinson's disease.

EP 0 900 568 A3



European Patent
Office

PARTIAL EUROPEAN SEARCH REPORT

Application Number

which under Rule 45 of the European Patent Convention EP 98 30 7181
shall be considered, for the purposes of subsequent
proceedings, as the European search report

DOCUMENTS CONSIDERED TO BE RELEVANT		
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim
A	WO 92 13535 A (DWIVEDI CHANDRADHAR ; OMODT GARY W. (US)) 20 August 1992 (1992-08-20) * page 2 - page 3; claim 1 *	1-7
A	US 4 183 931 A (RATHMAN TERRY L ET AL) 15 January 1980 (1980-01-15) * claims 1,2 *	1-7
A	KOE B K ET AL: "ENHANCEMENT OF BENZODIAZEPINE BINDING BY METHAQUALONE AND RELATED QUINAZOLINONES" DRUG DEVELOPMENT RESEARCH, US, NEW YORK, NY, vol. 7, no. 3, 1986, pages 255-268, XP002064447 ISSN: 0272-4391 * page 256 - page 257 *	1-7
A	WO 96 08493 A (NOVONORDISK AS ; NIELSEN FLEMMING ELMELUND (DK)) 21 March 1996 (1996-03-21) * page 5; claim 1 *	1-7
		<p>CLASSIFICATION OF THE APPLICATION (Int.Cl.6)</p> <p>A61K31/505 /(A61K31/505, 31:195,31:15)</p> <p>TECHNICAL FIELDS SEARCHED (Int.Cl.6)</p> <p>A61K A61P</p>
<p>INCOMPLETE SEARCH</p> <p>The Search Division considers that the present application, or one or more of its claims, does/do not comply with the EPC to such an extent that a meaningful search into the state of the art cannot be carried out, or can only be carried out partially, for these claims.</p> <p>Claims searched completely :</p> <p>Claims searched incompletely :</p> <p>Claims not searched :</p> <p>Reason for the limitation of the search:</p> <p>see sheet C</p>		
Place of search		Examiner
MUNICH		Brunnauer, H
Date of completion of the search		
8 February 2001		
<p>CATEGORY OF CITED DOCUMENTS</p> <p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p> <p>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document</p>		

EPO FORM 1503/03 R2 (P04C07)



European Patent
Office

INCOMPLETE SEARCH
SHEET C

Application Number
EP 98 30 7181

Claim(s) searched completely:
2-7

Claim(s) searched incompletely:
1

Reason for the limitation of the search:

Present claim 1 relates to an extremely large number of possible compounds. Support within the meaning of Article 84 EPC and/or disclosure within the meaning of Article 83 EPC is to be found, however, for only a very small proportion of the compounds claimed. In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search over the whole of the claimed scope is impossible. Consequently, the search has been carried out for those parts of the claims which appear to be supported and disclosed, namely those parts relating to the particular quinazoline-derivatives of claim 1.

PARTIAL EUROPEAN SEARCH REPORT

Application Number
EP 98 30 7181

DOCUMENTS CONSIDERED TO BE RELEVANT			CLASSIFICATION OF THE APPLICATION (Int.Cl.6)
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	
A	WO 95 21842 A (NOVONORDISK AS) 17 August 1995 (1995-08-17) * claims 1,9,10 *	1-7	
A	LOSCHMANN P. ET AL: "Synergism of the AMPA-Antagonist NBQX and the NMDA-Antagonist CPP with L-Dopa in Models of Parkinson's Disease." JOURNAL OF NEURAL TRANSMISSION. PARKINSONS DISEASE AND DEMENTIA SECTION., vol. 3, no. 3, 1991, pages 203-213, XP000973350 * page 205 * * page 207 * * page 211 - page 212 *	1-7	TECHNICAL FIELDS SEARCHED (Int.Cl.6)
A	WO 93 08173 A (SCHERING AG) 29 April 1993 (1993-04-29) * page 4 - page 6; claim 6 *	1-7	
D,A P,A	WO 97 43276 A (PFIZER ;ELLIOTT MARK LEONARD (US); WELCH WILLARD MCKOWAN JR (US)) 20 November 1997 (1997-11-20) * page 2, line 14,15 * * claims 1-22 *	1-7	

ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.

EP 98 30 7181

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report.
The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

08-02-2001

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9213535 A	20-08-1992	NONE	
US 4183931 A	15-01-1980	NONE	
WO 9608493 A	21-03-1996	AT 183186 I AU 694554 B AU 3380895 A CA 2200056 A CN 1193973 A CZ 9700753 A DE 69511420 D DE 69511420 T DK 781282 T EP 0781282 A ES 2138233 T FI 971076 A GR 3031788 T HU 76802 A JP 10506620 T NO 971187 A NZ 292108 A PL 319175 A SK 33497 A ZA 9507794 A	15-08-1999 23-07-1998 29-03-1996 21-03-1996 23-09-1998 13-08-1997 16-09-1999 30-03-2000 28-02-2000 02-07-1997 01-01-2000 13-05-1997 29-02-2000 28-11-1997 30-06-1998 15-05-1997 25-02-1999 04-08-1997 04-02-1998 14-05-1996
WO 9521842 A	17-08-1995	AU 692524 B AU 1805595 A CA 2182957 A CN 1145071 A,B EP 0743945 A FI 963130 A HU 76323 A IL 112588 A JP 9508632 T NO 963337 A NZ 281183 A US 5741785 A ZA 9501103 A	11-06-1998 29-08-1995 17-08-1995 12-03-1997 27-11-1996 09-10-1996 28-08-1997 14-07-1999 02-09-1997 11-10-1996 27-05-1998 21-04-1998 12-08-1996
WO 9308173 A	29-04-1993	DE 4135871 A AU 664212 B AU 2889492 A CA 2099270 A CN 1072929 A,B CZ 9301387 A EP 0565683 A	29-04-1993 09-11-1995 21-05-1993 27-04-1993 09-06-1993 19-01-1994 20-10-1993

EPO FORM P0453

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

ANNEX TO THE EUROPEAN SEARCH REPORT ON EUROPEAN PATENT APPLICATION NO.

EP 98 30 7181

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report.
The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

08-02-2001

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9308173 A		FI 932959 A	24-06-1993
		HU 64756 A	28-02-1994
		JP 6503583 T	21-04-1994
		NO 304693 B	01-02-1999
		NZ 244896 A	26-07-1995
		PL 299929 A	05-04-1994
		PT 101004 A,B	31-01-1994
		RU 2117663 C	20-08-1998
		SK 72793 A	06-10-1993
		ZA 9208275 A	04-05-1993
		MX 9303555 A	30-06-1994
WO 9743276 A	20-11-1997	AU 1554997 A	05-12-1997
		BG 102999 A	30-09-1999
		BR 9709085 A	03-08-1999
		CA 2252907 A	20-11-1997
		CN 1218464 A	02-06-1999
		CZ 9803526 A	17-11-1999
		EP 0901487 A	17-03-1999
		HR 970261 A	31-12-2000
		HU 9902148 A	28-04-2000
		JP 11514663 T	14-12-1999
		NO 985293 A	13-01-1999
		PL 330042 A	26-04-1999
		SK 152098 A	16-05-2000
		TR 9802296 T	22-02-1999
		ZA 9704156 A	16-11-1998

EPO FORM P0459

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82